

# **Scoped Environmental Impact Statement**

## **Petrie II Block 8**

### **Cumberland Ward, Ottawa, Ontario**

---

**Prepared for:**

Brigil Homes  
98, rue Lois  
Gatineau, Quebec  
J8Y 3R7

**Prepared by:**

Bowfin Environmental Consulting Inc.  
168 Montreal Road  
Cornwall, ON  
K6H 1B3

December 2015  
**(updated September 2021)**

## List of Acronyms and Definitions

ABBO - Atlas of Breeding Birds of Ontario  
ANSI – Area of Natural and Scientific Interest  
BHA - Butternut Health Assessment/Butternut Health Assessor  
BLTU – Blanding’s Turtle  
CC - Co-Efficient of Conservation  
CRZ - Critical Root Zone  
DBH - Diameter at breast height  
DFO – Fisheries and Oceans Canada  
EIS – Environmental Impact Statement  
ELC - Ecological Land Classification  
    CUM – Cultural Meadow  
ESA - *Endangered Species Act* (Provincial)  
GPS – Global Positioning System  
    NAD 83: North American Datum 1983  
    UTM: Universal Transverse Mercator  
LIO - Land Information Ontario  
MECP – Ministry of Environment, Conservation and Parks  
MNDMNR – Ministry of Northern Development, Mines, Natural Resources and Forestry  
MTO – Ministry of Transportation Ontario  
MVCA – Mississippi Valley Conservation Authority  
NHIC – Natural Heritage Information Centre  
NHRM - Natural Heritage Reference Manual  
OMNR/MNRF - Ontario Ministry of Natural Resources (old name)  
    -Ministry of Natural Resources and Forestry (old name)  
OP – Official Plan  
OWES - Ontario Wetland Evaluation System  
PPS - Provincial Policy Statement  
PSW - Provincially Significant Wetlands  
RVCA - Rideau Valley Conservation Authority  
SAR - Species at Risk (in this report they refer to species that are provincially or federally listed  
    as endangered or threatened and receive protection under ESA or SARA)  
SARA - *Species at Risk Act* (Federal)  
SARO - Species at Risk in Ontario  
SWH - Significant Wildlife Habitat  
SWHCS – Significant Wildlife Habitat Criteria Schedules for Ecoregion 6E  
SWHTG - Significant Wildlife Habitat Technical Guide

### **SRANK DEFINITIONS**

- S1** Critically Imperiled in the nation or state/province because of extreme rarity (often 5 or fewer occurrences) or because of some factor(s) such as very steep declines making it especially vulnerable to extirpation from the state/province.
- S2** Imperiled in the nation or state/province because of rarity due to very restricted range, very few populations (often 20 or fewer), steep declines, or other factors making it very vulnerable to extirpation from the nation or state/province.
- S3** Vulnerable in the nation or state/province due to a restricted range, relatively few populations (often 80 or fewer), recent and widespread declines, or other factors making it vulnerable to extirpation.
- S4** Apparently Secure; uncommon but not rare; some cause for long-term concern due to declines or other factors.
- S5** Secure; Common, widespread, and abundant in the nation or state/province.
- ?** Inexact Numeric Rank—Denotes inexact numeric rank

**SNR** Unranked, Nation or state/province conservation status not yet assessed.

**SU** Unrankable, currently unrankable due to lack of information or due to substantially conflicting information about status or trends.

**SNA** Not Applicable, A conservation status rank is not applicable because the species is not a suitable target for conservation activities.

**S#S#** Range Rank, A numeric range rank (e.g., S2S3) is used to indicate any range of uncertainty about the status of the species or community. Ranges cannot skip more than one rank (e.g., SU is used rather than S1S4).

**S#B** Breeding

**S#N** Non-Breeding

### **SARA STATUS DEFINITIONS**

**END** Endangered: a wildlife species facing imminent extirpation or extinction.

**THR** Threatened: a wildlife species that is likely to become endangered if nothing is done to reverse the factors leading to its extirpation or extinction.

**SC** Special Concern, a wildlife species that may become threatened or endangered because of a combination of biological characteristics and identified threats.

### **SARO STATUS DEFINITIONS**

**END** Endangered: A species facing imminent extinction or extirpation in Ontario which is a candidate for regulation under Ontario's ESA.

**THR** Threatened: A species that is at risk of becoming endangered in Ontario if limiting factors are not reversed.

**SC** Special concern: A species with characteristics that make it sensitive to human activities or natural events.

**Coefficient of Conservatism Ranking Criteria**

- 0 Obligate to ruderal areas.
- 1 Occurs more frequently in ruderal areas than natural areas.
- 2 Facultative to ruderal and natural areas.
- 3 Occurs less frequent in ruderal areas than natural areas.
- 4 Occurs much more frequently in natural areas than ruderal areas.
- 5 Obligate to natural areas (quality of area is low).
- 6 Weak affinity to high-quality natural areas.
- 7 Moderate affinity to high-quality natural areas.
- 8 High affinity to high-quality natural areas.
- 9 Very high affinity to high-quality natural areas.
- 10 Obligate to high-quality natural areas.

## Table of Contents

1.0	INTRODUCTION .....	7
2.0	METHODOLOGY .....	10
2.1	Background Review .....	10
2.2	Field Studies .....	10
2.2.1	Habitat Descriptions and Flora Observations .....	10
2.2.2	Butternut Inventory.....	11
2.2.3	Incidental Fauna Observations .....	11
3.0	Results .....	13
3.1	Background Review .....	13
3.2	Existing Conditions .....	19
3.2.1	Geology and Hydrologic Conditions.....	19
3.2.2	Vegetation Cover.....	22
3.2.3	Incidental Wildlife Observation .....	27
3.2.4	Aquatic Features .....	27
4.0	Potential to Impact the Natural Features .....	28
4.1	Impact Assessment Methods .....	31
4.2	Evaluation of Potential Impacts.....	32
4.2.1	Provincially Significant Wetlands.....	32
4.2.2	Natural Heritage System.....	34
4.2.3	Other – Urban Natural Area .....	37
4.2.4	SAR .....	39
4.2.5	Accidents and malfunctions.....	50
4.2.6	Other .....	50
5.0	CONCLUSIONS AND RECOMMENDATIONS .....	59
6.0	REFERENCES .....	60
	Appendix A : Background Review Mapping.....	63
	Appendix B: Review of Potential for Significant Wildlife Habitat.....	67
	Appendix C: SAR Hand-Out .....	74

**List of Figures**

Figure 1: General Location of Subject Lands ..... 9  
Figure 2: Location of the Study Area ..... 12  
Figure 3: City of Ottawa OP Schedule B ..... 17  
Figure 4: City of Ottawa OP Schedule L1 ..... 18  
Figure 5: Vegetation Communities ..... 21  
Figure 6: Site Plan with Vegetation Mapping ..... 29  
Figure 7: Spring View of Ravine ..... 30  
Figure 8: Potential Blanding's Turtle Habitat ..... 45

**List of Tables**

Table 1: Summary of Available Background Information on the Identified Natural Features (PSW, Woodlands, Valleylands, ANSIs, ESA, SWH, and Fish Habitat, and Policies under Section 2.4.2) ..... 15  
Table 2: Summary of Dates, Times, Conditions and Purpose of Site Investigations ..... 19  
Table 3: Summary of Soil and Geology Information Available from the Characterization of Ottawa's Watershed Maps ..... 20  
Table 4: Summary of Potential SAR ..... 40  
Table 5: Summary of Impacts, Mitigation Measures and Residual Effects ..... 52

**List of Photographs**

Photo 1: Cultural Meadow (CUM) (September 1, 2021) ..... 23  
Photo 2: Cultural Meadow (CUM) (September 1, 2021) ..... 23  
Photo 3: Fresh- Moist Bur Oak Deciduous Forest (September 1, 2021) ..... 25  
Photo 4: Narrow-leaved Emergent Marsh (September 1, 2021) ..... 25  
Photo 5: Park (September 22, 2015) ..... 26  
Photo 6: Looking north from the recreational path towards the Ottawa River at the ravine (September 22, 2015) ..... 35

## 1.0 INTRODUCTION

Bowfin Environmental Consulting Inc. (Bowfin) was retained by Brigil Homes, hereafter referred to as the proponent, to prepare a scoped Environmental Impact Statement (EIS) for Petrie's Landing Block 8 located at 180 Prestige Circle in support of their site plan application. The subject lands include approximately 0.7 hectares on the south side of the Jeanne d'Arc Boulevard North, approximately 1.1 km west of Trim Road, in part of Lot 33, Concession 1 of Cumberland Ward in the City of Ottawa (Figure 1). This original report was completed in 2015 and updated, with a site visit, in 2021.

It is Bowfin's understanding that the setbacks from the watercourse/ravine were already established at the Plan of Subdivision phase and that this current application follows those already approved setbacks.

It is noted that a Wetland Impact Study was completed by Muncaster Environmental Planning [*Wetland Impact Study for North Service Road Properties Cumberland Ward, City of Ottawa* (September 2004)]. That report also included a description of the terrestrial vegetative communities and a discussion of the potential impacts of development on the wetland, forests and other terrestrial areas and wildlife habitat.

Since the Muncaster Environmental Planning (MEP) 2004 report, portions of Phases 1 and 2 of Petrie's Landing have been built. Prior to constructing Phase 2 Block 8, the proponent was requested, by the City to update the 2004 report. To this effect Bowfin completed a site visit and an assessment of the natural environment to determine if any changes to the recommendations made in the above mentioned report were required. As per the Provincial Policy Statement (PPS) there are several natural features and areas identified as needing protection:

- Significant habitat of Endangered and Threatened Species;
- Significant wetlands;
- Significant woodlands;
- Significant valleylands;
- Significant wildlife habitat;
- Significant Areas of Natural and Scientific Interest; and
- Fish habitat.

The City of Ottawa protects these and other natural features through their Official Plan (OP) as described in the policies found in Section 2.4.2. All of the features listed in the PPS, but habitat of endangered or threatened species, are identified on the City's OP schedules as either Significant Wetlands, Natural Environment Area, Rural Natural Features, or Urban Natural Features (as applicable). For this project, the applicable schedules are: B, K and L1. The


presence/absence of habitat for endangered or threatened species are not depicted on the OP schedules. Instead, the appropriate provincial methodology [i.e. species-specific surveys, presence of preferred habitats] must be used to assess the potential impact to these species. The OP allows for other features, not identified on the schedules, that meet the criteria outlined in the *Natural Heritage Reference Manual* (NHRM) to be deemed significant through the EIS or other plans (i.e. CDP). If features are identified, then the potential to be negatively impacted is evaluated. The PPS states that a negative impact signifies:

*“a) in regard to policy 2.2, degradation to the quality and quantity of water, sensitive surface water features and sensitive ground water features, and their related hydrologic functions, due to single, multiple or successive development or site alteration activities;*  
*c) in regard to fish habitat, any permanent alteration to, or destruction of fish habitat, except where, in conjunction with the appropriate authorities, it has been authorized under the Fisheries Act;*  
*d) in regard to other natural heritage features and areas, degradation that threatens the health and integrity of the natural features or ecological functions for which an area is identified due to single, multiple or successive development or site alteration activities.”*

The following report provides a summary of the findings and an assessment of the functions and values of the natural features on the subject lands. It assesses the features to determine their significance following the applicable guidelines as referred to in the OP. The potential impacts to significant natural features are assessed and avoidance and mitigation measures provided.



Figure 1: General Location of Subject Lands



## 2.0 METHODOLOGY

Work undertaken for the completion of this project included a background review of existing information and field investigations.

### 2.1 Background Review

Where the OP indicated that the features to be considered were those identified on their schedules, these took precedent. Other information collected from outside sources was used to help inform the functions of these features and to identify those not found on the schedules (i.e. Endangered and Threatened species habitat). Outside sources included: Natural Heritage Information Centre (NHIC) database, iNaturalist, Atlas of Breeding Birds of Ontario (ABBO), Make-a-Map Land Information Ontario (LIO), and LIO databases. Information from personal knowledge has also been included as appropriate. The desktop review included a larger area (~5 km).

### 2.2 Field Studies

The initial field work completed in 2015, was update din 2018 and 2021.

#### 2.2.1 Habitat Descriptions and Flora Observations

Habitat mapping was completed through the use of satellite imaging and ground truthed during the field visits. The field studies were completed by systematically cruising the study area. Specific habitat types within the study area, identified during the preliminary mapping exercise were also targeted for community description. Habitat descriptions were based on the appropriate methodologies such as: *Ontario Wetland Evaluation System, Southern Manual* (OWES) for wetland habitats and the *Ecological Land Classification for Southern Ontario* (ELC) for terrestrial habitats. The MNR's ELC and OWES definition of wetlands do not match one another. Since wetlands are to be evaluated following OWES, the determination of the presence/absence of wetland habitat was based on the OWES definition of wetland habitat:

*“Lands that are seasonally or permanently flooded by shallow water as well as lands where the water table is close to the surface; in either case the presence of abundant water has caused the formation of hydric soils and has favored the dominance of either hydrophytic or water tolerant plants”.*

Specific attention was paid to locating species at risk (SAR) or species of conservation value listed as potentially occurring within the study area. If these species were observed, they would be photographed, and their coordinates recorded on a hand-held GPS using NAD83. Plants that could not be identified in the field were collected for a more detailed examination in the laboratory.

Nomenclature used in this report follows the Southern Ontario Plant List (Bradley, 2007) for both common and scientific names which are based on Newmaster *et al.* (1998). Authorities for scientific names are given in Newmaster *et al.* (1998).


### **2.2.2 Butternut Inventory**

Butternuts are an endangered species. While the Ministry of Environment, Conservation and Parks (MECP) is now responsible for the *Endangered Species Act* (ESA), they have not provided new guidelines. Previously, the MNRF certified Butternut Health Assessors (BHA) to complete Butternut Health Assessments as per MNRF's guidelines. This BHA was completed by a qualified Butternut Health Assessor (#281) in 2015. Presence of butternuts taller than the depth of snow was searched for on January 8, 2020 (BHA #723) and the inventory was repeated on September 1, 2021 (BHA #117). The search included the site and the adjacent 50 m around the site, to the south of Jeanne d'Arc. Any individuals noted would be marked with white spray paint and flagging tape and numbered sequentially. Their UTM's, using a GPS unit set at NAD83, would be recorded and the individual would be assessed according the BHA protocol.

### **2.2.3 Incidental Fauna Observations**

During the site visit any wildlife observations were recorded. Incidental observations included observations of an individual, its tracks, burrows, feces and/or kill sights.

Figure 2: Location of the Study Area



## 3.0 Results

### 3.1 Background Review


The subject lands, approximately 0.7 ha in size, are in Cumberland Ward of the City of Ottawa on Part of Lot 33 Concession 1 (Figure 1). They are situated to the north of Highway 174, west of Bellevue Ravine and south of Jeanne d'Arc Boulevard North Road. The lands to the west, east and south are developed. The designated land-use for the subject lands is General Urban Area on Schedule B of the City of Ottawa Official Plan (OP). The only natural heritage constraints those listed on Schedule L1 and are associated with the ravine to the east of the site, also referred to as Bellevue Ravine. This ravine is identified as a natural heritage system. North of the Jeanne d'Arc Boulevard North the habitat is natural, and the OP identified the following features: natural heritage system, urban natural feature, fish habitat (Ottawa River) and the Petrie Island Wetland. Petrie Island Wetland is a provincially significant wetland (PSW).

Muncaster (2004) cites McNeely (1995) as noting that Bellevue Ravine was not fish habitat and appeared to no longer convey significant flow, likely a result of upstream residential and road developments that redirected flow to either Brisbois Creek or Taylor Creek. Muncaster (2004) confirmed this during his August survey indicating that there was no defined watercourse among meadow marsh habitat in the Bellevue ravine south of the Jeanne d'Arc Boulevard North Road. This area was also noted as being dry or with moist soil with only a small area of minor ponding by Paterson Group on June 22, 2021 (Paterson Group Memo, June 23, 2021). It was noted that there was water present in the Ravine on September 1, 2021 but no defined channel in through the vegetation.

The City of Ottawa recently released its Significant Woodlands Guidelines. This Site is situated within the Urban Area and as per the Official Plan Section 2.4.2 iii, significant woodlands must meet both of the following conditions:

1. 0.8 ha in size (or larger)
2. Support woodland that is at least 60 years old at the time of evaluation. This threshold on age exceptions is depicted in Figure 2 of the Guidelines (included below).

**Figure 2. Application of the Size Threshold and Age Exemption in the Identification of Significant Urban Woodlands**



(from *Significant Woodlands: Guidelines for Identification, Evaluation, and Impact Assessment* (City of Ottawa, 2019))

A review of the 1975 image on geoOttawa depicts a narrower treed corridor than what is seen on site today. That image shows that the treed area along the ravine was at least 15 m from the existing property line. The air photo from 1965 indicates that there were even fewer trees on the west side of the ravine as compared to 1975 (Appendix A). Together this indicate that any wooded area on the property does not meet the minimum criteria. It is difficult to see the exact location of the trees in the 1965 air photo, regardless it is clear on the 1975 air photo on geoOttawa that any significant woodland is over 15 m from the current property line. It is important to note that this project has an approved Plan of Subdivision and the new guidelines for evaluating woodlands within the City of Ottawa Section 6.4.4.1 indicates that “...*new significant woodlands shall not be identified in those urban areas where the natural heritage system has already been identified in a current Secondary Plan, Community Design Plan, **Plan of Subdivision**, or an Existing Conditions Report submitted to and accepted by the City.*”

Table 1: Summary of Available Background Information on the Identified Natural Features (PSW, Woodlands, Valleylands, ANSIs, ESA, SWH, and Fish Habitat, and Policies under Section 2.4.2)

Natural Heritage Feature	Present within Site	Present within 120 m of Site	Additional Notes
Provincially Significant Wetlands (PSW)	No	The Petrie Island Wetland (PSW) is located 65m to the North	None
Habitats or species designated by ESA (Provincial)	Potential for endangered or threatened species needs to be determined following assessment of the suitable habitats in or near the site. Preliminary review of the satellite images suggest that there is a potential for bats and Butternuts in the adjacent lands. See section 5 of this report for more information.		None
Significant Woodlands	None	The 1975 air photo (geoOttawa) shows the trees to be at least 15m from the current property line. 1965 air photo suggests that trees that would meet the minimum age would be even further away.	Not applicable, the setbacks from the Plan of Subdivision have been adhered to (exemption under section 6.4.4.1 of the City's guidelines)
Unevaluated Wetlands	None	LIO shows unevaluated wetlands associated with significant woodlands and PSWs 55m north of subject lands	None
Significant Valleylands	No defined channel in the ravine.		None
Significant Wildlife Habitat (SWH)	None identified. Discussed in Section 5		None
Areas of Natural and Scientific Interest (ANSIs)	None		Schedule B and L1 do not identify ANSI.
Urban Natural Features	None	Schedule B shows an urban natural feature 15m north of subject lands	Not applicable, the setbacks from the Plan of Subdivision have been adhered to

Natural Heritage Feature	Present within Site	Present within 120 m of Site	Additional Notes
Forest Remnants, Corridors	None	The treed area along the ravine may be considered a forest remnant/natural corridor	Not applicable, the setbacks from the Plan of Subdivision have been adhered to
Groundwater features		None observed	None
Fish Habitat/Surface Water Features	None	The Ottawa River is situated roughly 85 m to the north. The Bellevue Ravine was found not to have a defined channel and was dry during site investigations by Muncaster. Paterson Group indicated ravine was dry or with moist soil apart from one small, ponded area in June of 2021. In September 2021, water was present in the ravine, but no defined channel was present in the vegetation.	none



Figure 3: City of Ottawa OP Schedule B





Figure 4: City of Ottawa OP Schedule L1



### 3.2 Existing Conditions

The site visits undertaken by Bowfin is provided in Table 2.

Table 2: Summary of Dates, Times, Conditions and Purpose of Site Investigations

Date	Time (h)	Staff	Air Temperature (Min-Max) °C	Cloud Cover (%) Beaufort Wind Scale [Descriptor (scale)]	Amount of rainfall 7 Days prior to Visit (mm)	Purpose
September 22, 2015	1300	S. St. Pierre	14.0 (7.4-22.6)	Clear Wind: light air (1)	8.6	Vegetation description, review of habitat in ravine and butternut inventory
January 8, 2020	n/a	C. Fontaine	-3 (-9.3 to 1.1)	n/a	n/a (winter conditions)	Tree inventory Butternut inventory
April 19, 2021	1300-1500	M. Lavictoire	21 (11.8-23.2)	Partial clouds Wind: Light Breeze (2)	25.8+	Butternut inventory Confirm edge of HWM, wetland Review of vegetation communities

C. Fontaine – Cody Fontaine – Fish and Wildlife Technologist

M. Lavictoire – Michelle (Nunas) Lavictoire – B. Sc. Wildlife Resources and M.Sc. Natural Resources

S. St. Pierre – Shaun St. Pierre – B. Sc. Biology

\*Min-Max Temp Taken From: Environment Canada. National Climate Data and Information Archive. Ottawa International Airport. Available <http://climate.weatheroffice.gc.ca/> [September 16, 2021]

+ Two significant rain events (August 28 and August 29, 2021 were both above 10 mm)

#### 3.2.1 Geology and Hydrologic Conditions

In general, the area was flat with the exception of the steep slopes of the Bellevue Ravine on the east side of the study area. No surface water or defined channels were present within ravine during the site visit (September 22, 2015). Paterson Group also noted that the ravine was dry or with moist soil with the exception of one ponded area (June 22, 2021). Surface water was present on September 1, 2021 but there had been two recent significant rain events (14.4 mm on August 28 and 10.4 mm on August 29, 2021) and there continued to be a lack of defined channel through the dense vegetation. The nearest other surface water feature was the Ottawa and Taylor Creek, both are outside of the subject lands. The Ottawa River is over 85 m to the north and Taylor Creek over 230 m to the east. There were no lakes, ponds, streams or groundwater seeps


on the property. The forested edge of the ravine is very steep and no areas that would serve as vernal pools were noted.

The area is identified as Clay Plains in the mapping from the *Characterization of Ottawa's Watershed: An Environment Foundation Document with Supporting Information Base* (March 2011). A summary of the information from the above mentioned report and maps is provided in Table 1. The soils map of the area shows the subject lands as having the Rideau soil association (which tends to have gray neutral heavy clay marine material) (*Soils of Regional Municipality of Ottawa-Carleton*).

Table 3: Summary of Soil and Geology Information Available from the Characterization of Ottawa's Watershed Maps

Map	Classification
Bedrock	Limestone and dolomite, interbedded
Surficial Geology	Glaciomarine, clay silt
Physiography Unit	Clay Plains
Permeability	Low
Overburden Depth	Shallow
Hydrological Soil Group	D

Figure 5: Vegetation Communities



### 3.2.2 Vegetation Cover

The 2015 findings indicated that the entire subject lands consisted of a Fresh-Moist Mixed Meadow. Portions of the site had been cleared previously and used for temporary staging during the construction of other phases. The adjacent lands to the south formed part of this same community. West of the site is now developed as part of other phases and also includes a parkland in the middle of the Prestige Circle. The forest along the ravine consisted of deciduous forest (fresh) that was classified as Fresh-Moist Bur Oak to the north and Dry-Fresh White Ash – Hardwood to the south. The vegetation communities were reviewed on September 1, 2021 and found to be similar with the following exceptions: the adjacent lands to the south were now developed (Block 7) and as such, the cultural meadow description was adjusted to that which remains and that there was little difference in the forest communities as such only the Fresh-Moist Bur Oak Deciduous Forest (FOD9-3) was kept as it was more representative. None of the communities identified are considered rare vegetation communities [*Significant Wildlife Habitat Technical Guide* (2000)]. A description of the subject lands and natural habitat to the northwest are provided below. Apart from the edge of the wetland habitat in the ravine, which was delineated with a hand-held GPS, the community boundaries are based on satellite image interpretation.

#### **Cultural Meadow (CUM)**

This community was present throughout the subject lands and continued southeast until the Highway 174. The community was highly disturbed with storage containers, discarded garbage, gravel pads and spoil piles. Outside of the bare soil, the dominant layer was the ground cover which was characterized by birds foot trefoil, Kentucky bluegrass, purple closer, common plantain, quack grass, New England aster, timothy, common ragweed, wild carrot, white sweet closer, burdock and reed canary grass.



Photo 1: Cultural Meadow (CUM) (September 1, 2021)



Photo 2: Cultural Meadow (CUM) (September 1, 2021)

**Fresh- Moist Bur Oak Deciduous Forest (FOD9-3)**

This deciduous community was found within the eastern side of the adjacent lands. It was composed of 95% tree cover which included 5% coniferous trees. In MEP's 2004 report the community was identified as being Dry-Fresh Poplar Ash Deciduous forest with a notable amount of bur oak present, but significant changes in the stands structures have occurred since 2004 most notably the death of the ash in the canopy layer. Such changes have altered the structure making bur oak the dominant tree present.

This forested community was present on a steep (45° slope) ravine. The canopy was 13-15 m tall and provided 40% canopy cover and contained lots of gaps, likely the result of the dead and dying ash trees. The dominant species was bur oak (95%, average 15 cm). The sub-canopy (5-10 m tall; 60% cover) was still strongly vegetated with bur oak followed by green ash, white birch, basswood and trembling aspen followed by American elm and red maple. There was more Manitoba maple and also the presence of muscledwood was noted on the southern side of this community. The understory (1-3 m tall; 30% cover) was variable but included: young green ash, Tartarian honeysuckle, prickly ash, purple flowering raspberry, common buckthorn, alternate leaved dogwood, and young black cherry. The ground layer (40% cover) included: large-leaved aster, late goldenrod, wild carrot, common strawberry, Virginia creeper, wild grape, poison ivy and northern lady fern.

The bottom of the ravine was vegetated with reed canary grass and could be classed as a narrow-leaved emergent marsh inclusion. Portions of this area also contained spotted jewel-weed, and New-England aster and red osier dogwood was present on the edge of the bank. This community was less than 0.2 ha in size with an abrupt transition to upland habitat.





Photo 3: Fresh- Moist Bur Oak Deciduous Forest (September 1, 2021)



Photo 4: Narrow-leaved Emergent Marsh (September 1, 2021)

### **Parkland**

The park consisted of remnant natural vegetation which was thinned out and supplemented with plantings and manicured grass. The main woody species in the natural area were: trembling aspen, American elm, white ash, Manitoba maple, choke cherry, and wild red raspberry. The herbaceous species in the natural area included: grasses, Canada goldenrod, and rough goldenrod. Examples of the plantings are: sugar maple, black maple, white oak, white spruce and white pine.



Photo 5: Park (September 22, 2015)

### ***Plant Species Discussion***

The plant species recorded were analyzed based for the following parameters: number of species, percent native, provincial rank (SRank), species at risk (Endangered or Threatened provincially) and co-efficient of conservation (CC). This analysis provides information on the level of disturbance to the site and special features.

A total of 74 species were identified of which 65% were native. This is above the percent non-native cover in most natural areas in southern Ontario (which usually has between 20-30% non-native cover Oldlam et al., 1995). The higher percentage of non-native plants can be attributed to the plant species documented on the subject lands which as a result of the recent land clearing and use as a temporary work area. This also affected the average coefficient of conservation (cc) value of 3.4 which also indicates an area with severely degraded conditions. [The CC provides

information on the species' tolerance to disturbance; those species with a high CC (maximum of 10) are highly sensitive].

All plants had a provincial SRank of S4, S5 or SNA signifying that the species recorded are apparently secure, uncommon but not rare (S4), secure, widespread and abundant in the nation or province (S5) or not applicable because the species is not a suitable target for conservation activities (i.e. non-native species) (SNA).

No Endangered, Threatened or species with a SRank of S3 or higher or listed as Special Concern, including no Butternuts, were found, in 2015, 2018 or 2021.

### **3.2.3 Incidental Wildlife Observation**

There were only a few incidental observations, these were: eastern chipmunk and grey squirrel. Both are common species

### **3.2.4 Aquatic Features**

As mentioned in the background review section Muncaster (2004) and McNeely (1995) indicated that Bellevue Ravine did not represent fish habitat. A review of the available satellite imaging and aerial photographs (geoOttawa) do not depict a continuous channel between Jeanne d'Arc Boulevard North and the Ottawa River (Figure 7).

On September 22, 2015, the lack of channel within the ravine between Jeanne d'Arc Boulevard North and Highway 174 was confirmed. During that visit, the ravine was dry (8.6 mm of rainfall was recorded at the Ottawa Airport within the 7 days preceding the site visit). Paterson Group's comments from their June 22, 2021 visit was that most of this area was dry or had moist soil with some minor ponding on the north end. On September 1, 2021, surface water was present but there continued to be no defined channel through the vegetation. The aquatic vegetation in the ravine bottom was composed of: reed canary grass, spotted jewel-weed, and New-England aster.

The top of the banks were well vegetated with herbaceous vegetation and woody species. The most common species were: large-leaved aster, common strawberry, common buckthorn, purple flowering raspberry, red osier dogwood, green ash, basswood, trembling aspen and bur oak (bur oak was the dominant live tree species).

#### 4.0 Potential to Impact the Natural Features

The development of Petrie II Block 8 will require the removal of the cultural meadow habitat within the subject lands (approximately 0.7 ha) and a few individual trees (see the Tree Conservation Report). The remnant forest that would meet the minimum age criteria from the City is at least 15 m from the edge of the property line and would not be impacted. Information provided by others noted the following:

1. As per the Rideau Valley Conservation Authority's comments to the project (June 9, 2021), the floodplain is located within the ravine (therefore below the top of slope).
2. Geotechnical limit is recommended to be 8 m from the top of slope (Paterson Group, June 23, 2021)
3. Bowfin confirmed that its original delineation of the edge of potential Blanding's Turtle and the high water mark (edge of area from which the Category 2 habitat's 30 m width is to be measured) in the April report was accurate on September 1, 2021. The potential Category 2 Blanding's Turtle habitat was established at 30 m from this delineated edge which also provides the 30 m setback (or more) from the high water mark.
4. The development will connect to the City's water and sanitary systems and the stormwater management water will be managed according to the Ministry of Environment, Conservation and Parks (MECP) regulations. No stormwater management ponds are predicted for this site.

The Rideau Valley Conservation Authority's comments (June 9, 2021) have indicated that original **Plan of Subdivision** determined the appropriate setback from the watercourse/ravine and that the proposed project meets those approved setbacks. Based on the above, the proposal meets the City's Policy 2 of Section 4.7.3.

Figure 6 shows the location of the top of slope, top of bank, 15 m from top of bank and the edge of the Blanding's Turtle Category 2 habitat which is also the same or more conservative representation of the 30 m from the high water mark. Apart from one tiny section, where the Blanding's Turtle Category 2 Habitat extends into the area to be fenced (70 m<sup>2</sup>), all grading will be outside of all of these limits.

The purpose of this report is to discuss if there are any changes to the mitigation measures outlined in the Muncaster (2004) report or any new measures required as a result of changes to the SAR. The potential to impacts these features, list of mitigation measures and a conclusion is provided below following the summary of the impact assessment methods.

Figure 6: Site Plan with Vegetation Mapping





Figure 7: Spring View of Ravine



#### **4.1 Impact Assessment Methods**

The assessment of the potential impacts is completed by analyzing the impact of various activities associated with the project. The development would include the following activities:

- Clearing of the disturbed meadow and removal of a few individual trees
- Grading and backfilling
- Construction of residences

The significance of the potential impacts is measured using four different criteria:

1. Area affected may be:
  - a. local in extent signifying that the impacts will be localized within the project area
  - b. regional signifying that the impacts may extend beyond the immediate project area.
2. Nature of Impact:
  - a. negative or positive
  - b. direct or indirect
3. Duration of the impact may be rated as:
  - a. short term (construction phase, 1-2 years)
  - b. medium term (3-4 years)
  - c. long term (>4 years).
  - d. permanent
4. Magnitude of the impact may be:
  - a. negligible signifying that the impact is not noticeable
  - b. minor signifying that the project's impacts are perceivable and require mitigation
  - c. moderate signifying that the project's impacts are perceivable and require mitigation as well as monitoring and/or compensation
  - d. major signifying that the project's impacts would destroy the environmental component within the project area.

## 4.2 Evaluation of Potential Impacts

### 4.2.1 Provincially Significant Wetlands

The Petrie Island PSW is situated to the north of the recreational bike path, north of the Jeanne d'Arc Boulevard North. It does not include any habitat within the subject lands. The nearest distance between the PSW and the subject lands is 110 m. **No changes from the discussion found in Muncaster (2004) are needed.** A summary of the Muncaster (2005) findings are provided below.

- Petrie Island Wetland will not be directly impacted as there is no wetland habitat on site. There will be a minimum distance of approximately 110 m of vegetated land between the north edge of development and the south edge of the PSW habitat.
- Potential indirect impacts on the wetland habitat are also minimized as all works will occur on the south of the Jeanne d'Arc Boulevard North and south of the other disturbances (a sanitary sewer was installed in the early 2000s along Jeanne d'Arc Boulevard North, and a recreational pathway).
- In addition to the above, the nearest wetland habitat to the north of the site is the insensitive meadow marsh dominated by reed canary grass. These habitats are tolerant of changes in water quality and other aspects of the moisture regime no such impacts are anticipated. The more sensitive features of Petrie Island, including the vegetation communities and species of interest are not in proximity to the site. The more ecologically significant areas are not found in the vicinity of the subject lands.
- It is noted that Bellevue Ravine provides a path for sediment-laden water to travel from the subject lands during construction to the wetland. The potential for poor water quality to leave the site can be mitigated through common best management practices listed below.
- The existing forested slopes adjacent to the development tablelands will be retained in its existing conditions, with a setback of 15 metres from the top-of-bank to the limit of development.
- The stormwater management will be designed to meet MECP's requirements and will consider the presence of the PWS on the north side of the Jeanne d'Arc Boulevard North and the potential for water leaving the site to reach the PSW via Bellevue Ravine.
- No new recreational pathways are proposed as part of this project and as such there is not potential for its development to increase human presence on the wetland habitat to the north.



**Conclusion:**

- The Petrie Island Wetland is a Provincially Significant Wetland however it is located 110 m from the proposed development of Petrie's Landing Phase II Block 8. Jeanne d'Arc Boulevard North and the recreational path separate the proposed development from the wetland.
- There is not potential for direct impacts to the PSW wetland.
- Indirect impacts could occur as a result of change in water supply or quality, sediment/erosion to the forested slope on Bellevue Ravine which drains to the wetland. Mitigation measures for this is provided below. Note that the overall hydrology of the PSW is controlled by the Ottawa River levels.

**Potential Impacts and Mitigation Measures:**

- Indirect impacts as a result of change in water supply or quality, erosion to the forested slope on Bellevue Ravine which drains to the wetland. These will be mitigated by:
  - Ensuring that the stormwater management plans meet MECP's requirements and take the presence of the ravine into account.
  - The protection of the vegetation within the 15 m setback from the top of slope.
  - During construction, an appropriate erosion and sediment control strategy will be developed, installed, monitored and maintained. This will include, at a minimum, the installation of sediment fence (countersunk) along the edge of the limit of development (along the edge of the forest). The proponent will undertake to monitor the construction process.
  - At this time, no trees on the top of the slope of Bellevue Ravine is forecasted. If this changes then a permit from the City will be required prior to removing trees greater than 10 cm in diameter (a Tree Conservation Report will address this separately). Note that there may be a few individual trees removed but the forested area is anticipated to be protected.
  - Any stock piles of soil or fill material would be stored at least 30 m from the top of slope and protected by silt fencing. The proponent will undertake to monitor the construction process.
  - Additional materials (*i.e.* rip rap, filter cloth and silt fencing) should be readily available in case they are needed promptly for erosion and/or sediment control.
  - Erosion and sediment control measures need to be maintained and will require daily inspection to ensure that they are working as intended. Additional inspections will be required after rainfall or storm events.
  - The sediment fencing would not be removed until the site is stable.
- No additional access to the wetland will be created (no trails).

<b>Area</b>	<b>Nature</b>	<b>Duration</b>	<b>Magnitude</b>
Local	Negative Indirect	Short to Medium Term depending on extent	Unlikely to occur (would occur as a result of an accident or malfunction resulting in sediment laden or contaminated water leaving the site)

#### **4.2.2 Natural Heritage System**

A natural heritage system was identified on Schedule L1. This feature consists of the Bellevue Ravine. Its attributes consisted of deciduous forests (Fresh-Moist Bur Oak Deciduous Forests), with a distinct ravine. The Tree Conservation Report (Bowfin, 2021) only identified two trees with a diameter at breast height larger than 50 cm within 20 m of the Site. The majority of the trees were <30 cm. This ravine did not have any defined channels and was dry during the August 2014 (Muncaster 2004), September 2015 visits, and mostly dry on June 22, 2021 (Paterson Group) but had some surface water on September 1, 2021. Two significant rain events occurred shortly before the September 1, 2021, visit and may explain the water present (August 28, 2021 – 14.4 mm and August 29, 2021 – 10.4 mm recorded at Ottawa Airport). Muncaster (2004) also referred to the McNeely (1995) that listed this area as not fish habitat. The feature does not meet the PPS, NHRM o definitions of significant: valleyland, PSW, ANSI and does not provide fish habitat. The setbacks were established during the **Plan of Subdivision** phase, and it is understood that this proposal meets all previous commitments.

The primary functions are limited to the protection of the ravine’s slope from erosion. Like all treed areas, it can also provide nutrient cycling, hydrological cycling, and clean air and long-term storage of carbon (NHRM 2005). While it may have historically provided a movement corridor, its value is now limited. There is no continuation of the ravine to the south of Highway 174 and while there are distinct connections between the PSW and the Taylor Creek ravine, to the east, there is not one to the Bellevue Ravine (Figure 7, Photo 6). Any value as habitat for endangered or threatened species and this is discussed further below. A review of the ravine and the SWHESC found that the ravine does not provide any significant wildlife habitat (Appendix B).



Photo 6: Looking north from the recreational path towards the Ottawa River at the ravine (September 22, 2015).

The PPS does not permit development in significant woodlands south and east of the Canadian Shield unless it has been demonstrated that there will be no negative impacts on the natural features or the ecological functions. A woodland is defined as a treed area, woodlot, or forested area. For the purposes of this report, a woodland included any community that was described as a treed swamp (deciduous, coniferous, or mixed), tall shrub or low shrub swamp composed of tree species, woodland or forest (regardless of tree size). The data was used in combination with satellite image interpretation to determine the size of the forest stands and the communities within the subject and adjacent lands were described using ELC.

The City of Ottawa recently released its Significant Woodlands Guidelines. This Site is situated within the Urban Area and as per the Official Plan Section 2.4.2 iii, significant woodlands must meet both of the following conditions:

3. 0.8 ha in size (or larger)
4. Support woodland that is at least 60 years old at the time of evaluation. This threshold on age exceptions is depicted in Figure 2 of the Guidelines (included below).

However as noted above, the City's guidelines states that where **Plan of Subdivisions** have been approved, no new significant woodlands will be designated.

**Conclusion:**

- The forested slope along the northwest side of the subject lands form part of the identified natural heritage system.
- The proposed development abuts the identified natural heritage system but respects the established setbacks from the **Plan of Subdivision** phase.
- The development does not propose any removal of the trees along the ravine's banks.
- This ravine is limited in its function. The primary function is to prevent erosion of the ravine slopes.

**Potential Impact and Mitigation Measures:**

- A minimum of 15 m setback from the top of slope has been established and prevents direct impact. This is to be clearly shown on the construction drawings and staked/surveyed on-site prior to clearing of vegetation.
- Indirect impacts could occur if the trees along the top of the slope are accidentally harmed resulting in less stability of the slope.
  - The removal of trees is not forecasted
  - Geotechnical advice was followed and the minimum setback from the top of slope exceeds their 8 m requirements.
- A permit from the City will be required prior to removing trees greater than 10 cm in diameter. See the Tree Conservation Report for more details.
- Sturdy fencing will be installed outside of the Critical Root Zone (CRZ) (defined by the City as 10 x the DBH) of the trunk of the trees to be retained.
  - No grading or activities that may cause soil compaction (such as heavy machinery and stockpiling of materials) will be allowed within the fenced area.
  - Furthermore, no machinery maintenance or refueling or stockpiling is permitted within 5 m of the outer edge of this fencing.
  - Exhaust fumes from all equipment will be directed away from the canopy of the trees to be retained.
  - If roots of trees to be retained become exposed during site alterations, they will be buried immediately with soil or covered with filter cloth or woodchips and kept moist until the roots can be buried permanently.
  - Any roots that must be cut will be cut cleanly to allow for healing.
  - At the request of the City, the fence will have signs posted every 20 m or so indicating that the purpose of the fence is to protect the vegetation on other side and is not to be removed until the construction is completed.
- Refer to the Tree Conservation Report for additional measures.

<b>Activity</b>	<b>Area</b>	<b>Nature</b>	<b>Duration</b>	<b>Magnitude</b>
Clearing of trees. This will be limited by the setback (min. 15 m from top of slope)	Local	Negative Indirect	Long Term to Permanent depending on extent	Provided that the vegetation within 15 m of the top of slope is protected, then no alterations to the function of the remnant forest/ravine/NHS are anticipated to occur. This setback follows that established during the <b>Plan of Subdivision</b> phase

#### 4.2.3 Other – Urban Natural Area

The Petrie Island Wetland also forms part of the Urban Natural Area #92: Petrie Islands and Mainland. This area has been described as a 288.2 ha parcel of alluvial islands, riparian deciduous swamp forests and mainland deciduous and mixed upland forests. UNA assessment of the area assigned high ranking for the UNA's:

- Connectivity
  - Connected to the Ottawa River and is adjacent to UNA 188 (Petrie West), UNA 93 (Taylor Creek Valley)
- Size and shape
  - Contains approximately 160 ha of interior habitat (primarily wetland habitat)
- Natural communities
  - High native flora [co-efficient of conservation (cc) 4.61 with 63 high-rate cc species]
  - Moderate to severe impacts from invasive species (including glossy buckthorn, common buckthorn and reed-canary grass all of which were found within the adjacent lands of this development proposal)
- Representative flora
  - Young to submature Green Ash Deciduous Swamp Forest
  - Submature United Maple, Silver Maple, Red Maple Deciduous Swamp Forest (dominant vegetation on alluvial islands)
  - Submature to mature Hackberry Deciduous Swamp Forest (small areas on all islands)
  - Deciduous Thicket Swamp
  - Reed canary grass Marsh
  - Cattail Marsh

- Shallow water aquatic associates
- Young to submature upland forest (Green Ash, White Birch and Red Maple – common in lower slopes and backshore)
- Mature upland Mixed Forest (Eastern Hemlock and Sugar Maple – small area of original Ottawa shore forest in Queenswood Forest)
- Sand barren (dune-like area on West Island)
- Significant flora and fauna
  - High level of native biodiversity
  - Faunal representation of both common urban breeding birds, herptiles and mammals
- Wildlife habitat
  - Large population of map turtles and Blanding’s turtles in wetlands and adjacent swamp forest, respectively
  - Provincially significant least bittern (SAR) and black tern (Special Concern), at least former breeding species, in open marsh habitat
  - Breeding habitat for Regionally significant raptor Cooper’s hawk in Queenswood Forest

**Conclusion:**

- This UNA consists of alluvial islands, riparian deciduous swamp forests and mainland deciduous and mixed upland forests
- The mapping for this UNA shows that, within the vicinity of the study area, it is restricted to the north side of Jeanne d’Arc Boulevard North.
- No direct impacts to this feature will occur.
- Potential for indirect impacts are restricted to the potential for sediment-laden water leaving the site via Bellevue Ravine. This has been dealt with in other sections.
- Note that the potential for Blanding’s Turtle is also dealt with in a separate section.

**Potential Impact and Mitigation Measures:**

- No additional mitigation measures for the UNA are required.

<b>Area</b>	<b>Nature</b>	<b>Duration</b>	<b>Magnitude</b>
Local	Negative Indirect	Long Term to Permanent depending on extent	Not anticipated to occur (would occur as a result of an accident or malfunction)

#### 4.2.4 SAR

Terrestrial and wetland Endangered and Threatened Species at Risk, on private land, are protected under provincial *Endangered Species Act*. It is noted that bird species protected under the *Species at Risk Act* (SARA) are protected by the *Migratory Bird Convention Act* (MBCA) on private lands. Within this report, the acronym SAR refers to only Endangered or Threatened species. No Special Concern species were identified and further they do not receive protection from ESA or SARA.

A list of potential SAR was compiled using various sources and identified up to roughly 5 km from the Site. The resulting list includes 12 potential SAR: 1 reptile (Blanding's turtle), 6 birds (least bittern, eastern whip-poor-will, chimney swift, barn swallow, bobolink, and eastern meadowlark), 4 mammals (little brown myotis, northern myotis, eastern small-footed myotis, and the tri-colored bat), and 1 plant (butternut) (Table 4). Fish were not included as there is no fish habitat present. Of these, many were determined not to be present or had no triggers for review based on guidance from the province. Table 4 notes the relevant provincial guidelines and triggers and indicates whether the species is brought forward for discussion.

Table 4: Summary of Potential SAR

Common Name/ Population	Scientific Name	SRank	ESA Reg. 230/08 SARO List Status	SARA Schedule 1 List of Wildlife SAR Status	Preferred Habitat	Reference	MECP Guidelines/Triggers for Review	Brought Forward (Yes/No)
<b>REPTILES</b>								
Blanding's Turtle	<i>Emydoidea blandingii</i>	S3	THR	THR	Shallow water, large marshes, shallow lakes or similar such water bodies.	COSEWIC 2016a	This species is noted in the background information to be present within 500 m and the bottom of the ravine could provide habitat.	yes
<b>BIRDS</b>								
Least Bittern	<i>Ixobrychus exilis</i>	S4B	THR	THR	Freshwater marshes, ditches, creeks, rivers and lakes with tall emergent vegetation.	COSEWIC 2009	No habitat is present on-site, but individuals have been listed to occur within the PSW to the north of Jeanne d'Arc Boulevard North. MECP does not have category guidelines for this species and the wetland will not be impacted.	No
Eastern Whip-poor-will	<i>Caprimulgus vociferus</i>	S4B	THR	THR	Rock or sand barrens with scattered trees, savannahs, old burns or other disturbed sites in a state of early to mid-forest succession, or open conifer plantations	COSEWIC 2009	No suitable habitat is present on-site or within 500 m	No
Chimney Swift	<i>Chaetura pelagica</i>	S4B, S4N	THR	THR	Cities, towns, villages, rural, and wooded areas.	COSEWIC 2007	None observed, and most trees were <30 cm in diameter. No individuals are shown within 2 km of the site on iNaturalist, but they are present within 10 km (ABBO)	Yes
Barn Swallow	<i>Hirundo rustica</i>	S4B	THR	THR	Open or semi-open lands: farms, field, marshes.	COSEWIC 2011a	There were no structures in or within 5 m of the Site. Houses and buildings were present within 200 m, but these would not be impacted.	No
Bobolink	<i>Dolichonyx oryzivorus</i>	S4B	THR	THR	Primarily in forage crops, and grassland habitat.	COSEWIC 2010	Meadows are broadleaf, smaller than 4 ha, and not suitable grassland.	No



Common Name/ Population	Scientific Name	SRank	ESA Reg. 230/08 SARO List Status	SARA Schedule 1 List of Wildlife SAR Status	Preferred Habitat	Reference	MECP Guidelines/Triggers for Review	Brought Forward (Yes/No)
Eastern Meadowlark	<i>Sturnella magna</i>	S4B	THR	THR	Fields, meadows and prairies.	COSEWIC 2011b	Meadows are broadleaf, smaller than 4 ha, and not suitable grassland.	No
<b>MAMMALS</b>								
Little Brown Myotis	<i>Myotis lucifugus</i>	S4	END	END	Buildings, attics, roof crevices and loose bark on trees or under bridges. Always roost near waterbodies.	Eder 2002	MECP recommends the use of avoidance timing window for clearing of trees (>10 cm in diameter) if this can be accomplished then no impacts.	Yes
Northern Myotis/Northern Long-eared Bat	<i>Myotis septentrionalis</i>	S3	END	END	Older (late successional or primary forests) with large interior habitat.	Menzel et al. 2002, Broders et al. 2006, SWH 6E Ecoregion Criterion Schedule		
Eastern Small-footed Myotis	<i>Myotis leibii</i>	S2S3	END	No Status	Found within deciduous or coniferous forests in hilly areas.	Eder 2002		
Tri-colored Bat	<i>Perimyotis subflavus</i>	S3?	END	END	Prefers shrub habitat or open woodland near water.	Eder 2002		
<b>PLANTS</b>								
Butternut	<i>Juglans cinerea</i>	S3?	END	END	Variety of sites, grows best on well-drained fertile soils in shallow valleys and on gradual slopes	COSEWIC 2003	Inventory completed in 2015 and none found. No large individuals were noted during winter 2018. Inventory repeated on September 1, 2021. No butternuts were found. Inventory has a shelf-life of two-years (until September 2, 2023)	Yes

Status updated: March 2021

**SRANK DEFINITIONS**

**S1** Critically Imperiled, Critically imperiled in the nation or state/province because of extreme rarity (often 5 or fewer occurrences) or because of some factor(s) such as very steep declines making it especially vulnerable to extirpation from the state/province.

- S2** Imperiled, Imperiled in the nation or state/province because of rarity due to very restricted range, very few populations (often 20 or fewer), steep declines, or other factors making it very vulnerable to extirpation from the nation or state/province.
- S3** Vulnerable, Vulnerable in the nation or state/province due to a restricted range, relatively few populations (often 80 or fewer), recent and widespread declines, or other factors making it vulnerable to extirpation.
- S4** Apparently Secure, Uncommon but not rare; some cause for long-term concern due to declines or other factors.
- S#S#** Range Rank, A numeric range rank (e.g., S2S3) is used to indicate any range of uncertainty about the status of the species or community. Ranges cannot skip more than one rank (e.g., SU is used rather than S1S4).
- ?** Inexact Numeric Rank—Denotes inexact numeric rank
- S#B** Breeding
- S#N** Non-Breeding

**SARO STATUS DEFINITIONS**

- END** Endangered: A species facing imminent extinction or extirpation in Ontario which is a candidate for regulation under Ontario's ESA.
- THR** Threatened: A species that is at risk of becoming endangered in Ontario if limiting factors are not reversed.
- SC** Special Concern: A species with characteristics that make it sensitive to human activities or natural events.

**SARA STATUS DEFINITIONS**

- END** Endangered, a wildlife species facing imminent extirpation or extinction.
- THR** Threatened, a wildlife species that is likely to become endangered if nothing is done to reverse the factors leading to its extirpation or extinction.
- SC** Special Concern, a wildlife species that may become threatened or endangered because of a combination of biological characteristics and identified threats.

## Reptiles

### Blanding's Turtle

Blanding's turtle is associated with a variety of shallow slow aquatic habitats with submergent and emergent plants. These turtles require basking sites located near the water such as exposed rocks or partially submerged logs. The nesting sites are located within areas of loose substrates varying from sand to cobblestone and may occur along roadways as far as 400 m away. Marsh habitat is important for the juveniles for protection from predators. The species overwinters within permanent water bodies (COSEWIC, 2005). This species can migrate far distances of up to 6 km (OMNR, 2013b). Migration routes can include overland movement.

The habitat guidelines for Blanding's turtle provide protection to the areas surrounding a nest, or perceived nest area. The level of protection varies with the distance from the nest and has been categorized by MNRF into three categories. These, along with their protection level are:

- Category 1 Nest and the area within 30 m or Overwintering sites and the area within 30 m
- Category 2 The wetland complex (i.e., all suitable wetlands or waterbodies within 500 m of each other) that extends up to 2 km from an occurrence, and the area within 30 m around those suitable wetlands or waterbodies
- Category 3 Area between 30 m and 250 m around suitable wetlands/waterbodies identified in Category 2, within 2 km of an occurrence

No Blanding's Turtle surveys were undertaken. The habitat on-site did not provide overwintering, nesting or suitable movement corridor functions. There are occurrences of Blanding's turtle on the Ottawa River within the PSW (make-a-map) but no sightings on iNaturalist despite this area being heavily visited by the public. Blanding's Turtle are anticipated to utilize the aquatic habitat adjacent to the site (Ottawa River and associated wetland habitat). The Ottawa River could provide overwintering habitat.

The guidelines indicate that suitable habitat within 500 m of the sighting plus its adjacent habitat (30 m) should be automatically considered Category 2 habitat. The exact location of the sightings are unknown. It is assumed that the reed canary / jewelweed inclusion at the bottom of the ravine could provide suitable habitat for this species, and this is within 500 m of the PSW. While the Category 3 habitat would also be automatically placed over much of the site, the surrounding lands are cleared and developed and as not suitable for use. Based on these investigations, it is proposed that:


1. The Ottawa River be considered appropriate habitat for overwintering or Category 2.

2. The reed inclusion community at the base of the ravine be considered appropriate habitat for Category 2.

The value of much of the adjacent lands as Category 3 Habitat is questionable. The purpose of the Category 3 Habitat is to provide a migration corridor. To be suitable habitat, it should link wetland habitats or nesting habitats or overwintering areas. The surrounding areas to the west, east and south are developed. No overwintering, wetland or nesting areas are noted in these directions. The more natural migration route would be for the turtles to travel to the stormwater management pond to the west for overwintering or to migrate through Taylor Creek.

The development of Petrie II Block 8 would not affect the use of any existing habitat or migration routes. The mitigation measures would include the installation of a permanent exclusion fence (Figure 8).

Figure 8: Potential Blanding's Turtle Habitat



## *Birds*

### *Chimney Swift (Chaetura pelagica)*

The chimney swift can often be found in developed areas and prefers to utilize structures such as large (>50 cm diameter) trees or man-made structures such as chimneys for its nesting habitat (COSEWIC, 2007). The use of large trees is now considered a rare event and the documented occurrences have all be in trees that were <1 km from a waterbody (large enough to be shown on 1:50,000 topographical maps) (COSEWIC, 2007).

The results from the Tree Conservation Report were reviewed and there were only two trees that were suitable in size (diameter >50 cm) nearby. Neither will be impacted by this project. This species is easily identified when present, it is very vocal and forages often. There are no recordings of this species within 2 km of this site on iNaturalist and there are no large trees to be removed. While there remains a potential for its presence, impacts to this species can be avoided through the implementation of timing windows.

### *Barn Swallow (Hirundo rustica)*

The barn swallow can often be found nesting on man-made structures. The *General Habitat Description for Barn Swallow* (OMNRF, 2018b) indicates that the protected habitat for this species includes three categories:

- Category 1    nest
- Category 2    the area within 5 m of the nest
- Category 3    the area between 5 m and 200 m of the nest

No buildings were present within the subject lands. The buildings in the adjacent lands could provide habitat for this species, these will not be impacted by the developments. This species will not be impacted.

## *Bats*

The potential SAR bats within the general area are: little brown myotis, northern myotis, eastern small-footed myotis and tri-colored bat. There are three types of habitats required by bats: hibernation, maternity sites and day-roost sites. The latter is not considered critical habitat. These four bat species prefer to hibernate in caves or mines. They can hibernate in buildings but that is rare for these species (COSEWIC, 2013a). No caves, buildings, or mines were present.

The northern myotis tends to prefer larger expanses of older forests (late successional or primary forests) and chose maternity sites in snags that are in the mid-stage of decay. They prefer habitat with intact interior habitat and is shown to be negatively correlated with edge habitat (Menzel et al., 2002; Broders et al., 2006; Yates et al., 2006; OMNRF, 2015). This habitat is absent.

The recovery strategy for the eastern small-footed myotis indicates that the preferred maternity habitat of this species consists of open rock habitats and that it rarely uses old buildings as roosting/maternity sites (Humphrey, 2017). There was no suitable rocky habitat present or buildings. Based on this information, this species' maternity sites are considered absent.

The Atlas of Mammals of Ontario (Dobbyn, 1994) suggests that the tri-colored bat is not present within this part of Ontario however, the NatureServe mapping in the COSSARO (2015) includes all of southeastern Ontario. Based on this information, this species is considered to have a very low potential of occurring.

This leaves only the little brown myotis as potentially using the study area for maternity sites. There remains the potential for the other species to utilise the trees on-site for day-roosts. Mitigation measures will be included discussed further below.

## *Plants*

### *Butternuts*

As discussed above, no butternuts were identified in or within 50 m of this site by the surveyor in 2015 and no large trees were noted in the winter of 2020. Butternut inventory was completed on September 1, 2021, and still no butternuts were found. This species is considered absent but has been brought forward for discussion because butternut inventories have a shelf-life of 2-years. The last inventory is valid until September 1, 2023 (inclusive). If clearing of vegetation has not been completed by that time, then a new butternut inventory would be needed within 2-years prior to clearing of vegetation.

### **Mitigation Measures:**

#### General:

- Endangered and Threatened species are protected and cannot be harmed, harassed, or killed and in some cases their habitats are also protected. These individuals will only be handled by qualified person and only if the individual is in imminent threat of harm. An authorization under the ESA 2007 would be required to handle individuals that are not in imminent threat of harm.
- If a SAR enters the work area during the construction period, any work that may harm the individual is to stop immediately and the supervisor will be contacted. No work will

continue until the individual has left the area. These sightings will be reported to MECP and NHIC.

- Should an individual be harmed or killed then work will stop and MECP will be contacted immediately.

### Turtles

- Based on the mapping there is very little (70 m<sup>2</sup>) Category 2 lands that will be impacted (temporarily or permanently).
- During construction, an exclusion fences will be in place. The sediment fencing along the banks can be used for temporary exclusion fencing. These will be properly countersunk and maintained to ensure that any turtles cannot get into the site. This sediment fencing is, at a minimum, to include the side closest to the ravine. Reptile and Amphibian Exclusion Fencing: Best Practices (OMNR, 2013d) should be followed for exclusion fence design.
- A permanent barrier to turtle access of the newly developed area will be included in the final design of the development. The location has been shown on Figure 8. Reptile and Amphibian Exclusion Fencing: Best Practices (OMNR, 2013d) will be followed for exclusion fence design.
- Implement a strict speed limit of <15 km/h during construction.
- If possible, clearing of vegetation will take place outside of the active turtle season [i.e. clear after October 16 (or freeze up) and before April 15 (or spring thaw)].
- If clearing takes place during the active season, then a biologist familiar with this species, will sweep the area to be cleared immediately prior to the clearing and remain on site during clearing works. Where feasible clear using hand tools/chain saws.
- During clearing of vegetation, contractors are to be informed that they should keep a look out for wildlife and if any are observed, they should be given the opportunity to leave the area.
- Recommend clearing from west to east direction to allow wildlife the opportunity to leave the site into the natural areas that are to remain.
- Stockpiles that might provide suitable nesting substrate (i.e. gravel, soil) will be provided with additional sediment fencing to prevent turtles from nesting in the work area. Note that should Blanding's Turtle nest on-site, then all work would be stopped until the hatchlings leave in the fall and MECP would need to be contacted.
- Contractor is to perform daily sweeps during the active season (approximately April 15 to October 16, subject to weather conditions).
- If an individual is found, work that puts the individual in danger will cease (i.e. moving machinery), and the individual will be watched from far to document where and when it leaves the site for a minimum of 2 hours. If it does not leave, then it may need to be relocated. Contact a biologist experienced with this species to relocate the individual.



<b>Activity</b>	<b>Area</b>	<b>Nature</b>	<b>Duration</b>	<b>Magnitude</b>	<b>Likelihood</b>
Removal of vegetation	Local	Negative Direct	Permanent	Low potential; all work is outside of Category 2 habitat	Negligible (if timing window is followed and exclusion fences installed)
					Confirmation from MECP is pending

**Birds**

Apart from the low potential for chimney swifts to utilize the larger diameter trees in the ravine, there is not potential for critical SAR birds in the Site or adjacent lands.

**SAR Birds:**

- No trees that have a diameter of 50 cm or larger will be removed.
- The removal of all trees will occur outside of the Chimney Swift nesting period (provided from MECP as being between May 15 and August 31 in Southern Ontario (including eastern Ontario)). This is to ensure no disturbance to any that may be nesting in the adjacent lands. To remove this condition, two breeding bird visits would be required during the appropriate timing window (end of May to first week in July and spaced at least 15 days apart). However, it is noted that the bat timing window includes this period and as such, it cannot be removed without a bat exit survey as well (see below).

<b>Activity</b>	<b>Area</b>	<b>Nature</b>	<b>Duration</b>	<b>Magnitude</b>	<b>Likelihood</b>
Removal of vegetation	Local	Negative Direct	Permanent	Low potential; no critical SAR bird habitat within the area to be cleared	Negligible (if timing window is followed)

**Bats:** Recent discussions with MECP on bats, in the Kemptville area, indicate that they do not need to be approached if the timing window below can be adhered to.

- Educate contractors by informing them that most bats in Ontario are protected.
- Remove trees between October 1 and March 31 (Bat active season is currently assumed to be April 1 to September 30). If this is not possible, conduct exit survey prior to cutting them down. If the exit survey identifies bats, contact MECP or biologist for additional guidance.

<b>Activity</b>	<b>Area</b>	<b>Nature</b>	<b>Duration</b>	<b>Magnitude</b>	<b>Likelihood</b>
Removal of vegetation	Local	Negative Direct	Permanent	Low potential; habitat is not a restrictive item in eastern Ontario	Negligible (if timing window is followed and exclusion fences installed)

#### **4.2.5 Accidents and malfunctions**

The potential impacts associated with this proposed development largely stem from accidents or malfunctions. Although the likelihood of accidents and malfunctions occurring would be minimized by following the mitigation measures outlined below, should accidents and/or malfunctions occur they have the possibility of presenting serious impacts and require consideration.

Maintenance on construction equipment such as refueling, oil changes or lubrication would only be permitted in designated area located at a minimum of 30 m from the slope and in an area where sediment erosion control measures and all precautions have been made to prevent oil, grease, antifreeze or other materials from inadvertently entering the ground or the surface water flow.

Machinery should be cleaned prior to arriving on-site to prevent the potential spread of invasive species.

Emergency spill kits would be located on site. The crew would be fully trained on the use of clean-up materials in order to minimize impacts of any accidental spills. The area would be monitored for leakage and in the unlikely event of a minor spillage the project manager would halt the activity and corrective measures would be implemented. Any spills would be immediately reported to the MECP Spills Action Centre (1800 268-6060).

#### **4.2.6 Other**

The measures outlined above serve to protect the identified or potentially present natural features identified in the background review and/or site investigations. However, there are also some other items that should be mentioned.

1. Almost all birds in Ontario are protected by either MBCA or FWCA.
2. Most reptiles are protected by the FWCA

**Mitigation Measures:**

- Almost all breeding birds are protected under the MBCA and/or FWCA. The only species not protected are: American crow, brown-headed cowbird, common grackle, house sparrow, red-winged blackbird, and starling. It is prohibited to destroy or disturb an active nest of other birds, or to take or handle nests, eggs, or nestlings. In this part of Ontario, the current standard nesting period is between April 5<sup>th</sup> to August 28<sup>th</sup>. Outside of this timing window, it is considered unlikely that birds would be nesting. Note, there are some birds (birds of prey, herons etc.) that do begin nesting earlier in the year. It should also be noted, that if an active nest is present before or after the above dates that it is still protected. These dates only serve as a guideline.
- During construction, there is a potential for suitable habitat for ground nesting birds (i.e. killdeer) to be created. These include bare soil or gravel areas. Perform regular walks of the cleared areas looking for ground nesters. If any are present, the contact a biologist for guidance.
- Work during the daytime hours to prevent light disturbances.
- Ensure that all equipment have the appropriate mufflers to reduce noise disturbances.
- If a turtle nest is suspected, then flag a 10 m buffer to protect the nest. Contact MECP (for SAR) and MNR (all other species).

Table 5: Summary of Impacts, Mitigation Measures and Residual Effects

Activity	Natural Heritage Feature/Function	Potential Effect	Proposed Mitigation	Residual Effect
<b>Construction</b>				
Vegetation Clearing in preparation development	<p>Ravine</p> <p>Small potential for Blanding’s Turtle, Chimney Swifts and bats in adjacent lands.</p> <p>Bird nests protected by MBCA</p>	<p>Disturbance to vegetation along the slope of the ravine could result in erosion. Sediment-laden water could end up in the PSW downstream.</p> <p>If Blanding’s Turtle are present, they could be accidentally harmed by machinery if present during clearing.</p> <p>70 m<sup>2</sup> of Category 2 Blanding’s Turtle habitat would be lost.</p> <p>If Chimney Swifts are present in adjacent lands, they could be indirectly impacted by removal of vegetation</p>	<p>Vegetation within 15 m from the top of slope will not be impacted. This respects the <b>Plan of Subdivision</b> commitments.</p> <p>Only a few (2 live) trees on the site are anticipated to be removed. A permit from the City will be required prior to removing trees greater than 10 cm DBH. Refer to the Tree Conservation Report for details.</p> <p>Use small machinery within 20 m of ravine.</p> <p>Any clearing of vegetation within the</p>	None

Activity	Natural Heritage Feature/Function	Potential Effect	Proposed Mitigation	Residual Effect
		<p>during their nesting period.</p> <p>If bats are present, they could be impacted if trees (&gt;10 cm) are removed during their active season.</p> <p>Birds in general could be using the area for nesting.</p>	<p>CRZ (DBH x 10 cm) of trees to be retained will be done by hand tools.</p> <p>All vegetation clearing must occur outside of breeding bird season, active turtle season, active bat season.</p> <p>Exceptions can only be made if Chimney Swift surveys and bat exit surveys are completed along with general nesting bird surveys.</p> <p>Daily sweeps for turtles will be completed between April 15-October 16.</p> <p>If clearing takes place during active turtle season, a biologist will complete the daily sweeps.</p>	

Activity	Natural Heritage Feature/Function	Potential Effect	Proposed Mitigation	Residual Effect
			<p>Temporary (during construction) and permanent turtle exclusion (operation) will be installed and monitored.</p> <p>Stockpiles will have additional sediment fence around them to ensure that turtles do not nest.</p>	
Grading	Indirect impacts to wetland, ravine and UNA should erosion or sediment control measures fail.	Negative impacts to: quality of wetland or UNA habitat or its functions (wildlife and fish habitat) as a result of erosion or sedimentation of wetlands or aquatic habitats. Given the distance between the site and the natural features it is unlikely that even indirect impacts will occur to	Install erosion and sediment protection measures prior to the removal of vegetation. Erosion and sediment protection measures will include at a minimum properly keyed in sediment fencing (the heavy duty geotextile fabric needs to be buried to prevent water from traveling under the fence) along the top of	None provided that mitigation measures are properly implemented and maintained.

Activity	Natural Heritage Feature/Function	Potential Effect	Proposed Mitigation	Residual Effect
		<p>the Petrie Island PSW or UNA.</p> <p>Noise from machinery may also cause a disturbance to wildlife in the ravine.</p> <p>Permanent structures could cause slope instability.</p>	<p>slope of the ravine and around spoil piles.</p> <p>Maintain sediment fencing as needed.</p> <p>Daily inspections, especially following rain or storm events, of the sediment control measures will be required.</p> <p>Leave erosion control measures in place until slope is fully stabilized.</p> <p>No work outside of limit of development.</p> <p>No maintenance of equipment or fueling within 30 m of the ravine (this matches the edge of the grading area).</p>	

Activity	Natural Heritage Feature/Function	Potential Effect	Proposed Mitigation	Residual Effect
			<p>No storage of stockpiles within 30m of top of ravine (slope).</p> <p>Work during the daytime hours to prevent light disturbances.</p> <p>Ensure that all equipment have the appropriate mufflers to reduce noise disturbances.</p> <p>Slope stability to be confirmed by a geotechnical expert as needed.</p> <p>Construction staff will be informed of the SAR in the area (Appendix C).</p> <p>Proponent will conduct monitoring to confirm</p>	



Activity	Natural Heritage Feature/Function	Potential Effect	Proposed Mitigation	Residual Effect
			that contractors are following measures.	
Accidents or Malfunctions	Indirect impacts to wetland, ravine or UNA should erosion or sediment control measures fail.	Spills or accidents during construction could impact the quality of wetland or UNA or ravine habitats or their functions (wildlife and fish habitat).	<p>All equipment should be well maintained, clean and free of leaks.</p> <p>Maintenance of construction equipment should occur at a minimum of 30m from the top of the slope/ravine and in an area where all precautions have been made to prevent oil, grease, antifreeze or other materials from inadvertently entering the ground or surface water.</p> <p>Any machine coming from offsite should be cleaned and free of mud (to prevent the transfer of non-native vegetation).</p>	Unlikely

Activity	Natural Heritage Feature/Function	Potential Effect	Proposed Mitigation	Residual Effect
			<p>Emergency spill kits should be located on site and the crew trained on their use.</p> <p>Any spills will be reported immediately to MECP Spills Action Centre (1.800.268.6060).</p>	

## 5.0 CONCLUSIONS AND RECOMMENDATIONS

The subject lands consisted of a disturbed meadow. Nearby natural habitats consisted of the forested ravine, referred to as Bellevue Ravine. The meadow was disturbed during construction of other phases and roadways. The area is bordered by Jeanne d'Arc Boulevard North to the north, Highway 174 on the south and surrounded by other development on the east and west. The natural habitat north of the Jeanne d'Arc Boulevard North and recreational bike path consisted of Petrie Island PSW, Ottawa River and an identified natural heritage system (UNA #92). These significant features will not be directly impacted by the proposed development. They could be indirectly impacted if a large sediment or contaminant spill occurred during construction however given the project's location and distance from these it is considered unlikely especially if properly installed and maintained sediment and erosion control practices are followed.

The setback requirements for this property were established during the **Plan of Subdivision** phase. The RVCA comments noted that those approved setback conditions were met with this proposed phase. Since that time, the potential for Blanding's Turtle became a consideration. The existing plan is able to meet the typical requirements for MECP approval. The potential loss of the small (70 m<sup>2</sup>) amount of Category 2 habitat and the measures proposed herein are being reviewed by MECP. The Bellevue Ravine and its slopes are to be protected through the minimum setback of 15 m from top of slope. This will protect the remnant forest associated with the ravine. This setback exceeds the 8 m geotechnical setback.


No SAR habitat or species were documented on the subject lands. No raptor nests were found within this area. Butternuts were confirmed to be absent on September 1, 2021. With respect to species at risk, the most likely species would be: Blanding's turtle, chimney swift or bats. Avoidance and mitigation measures have been included to prevent harm to these or their habitats. It will be circulated to MECP for their review.

All of the impacts can be mitigated through the use of common mitigation measures and no residual negative impacts to the natural environment are anticipated as a result of the development of Phase II Block B. Unless, MECP provides additional measures, this proposed development can be accepted as planned.

I trust that this report will meet your requirements. Should you have any questions or comments, please contact the undersigned.

Sincerely,

Bowfin Environmental Consulting Inc.



---

Michelle Lavictoire, Biologist / Principal

## 6.0 REFERENCES

- Bradley, David. 2007. Southern Ontario Vascular Plant Species List. Prepared by Southern Science and Information Section, Ontario Ministry of Natural Resources, Peterborough, Ontario. 57pp.
- Broders, H., Forbes, G., Woodley, S. & Thompson, I. (2006). Range extent and stand selection for roosting and foraging in forest-dwelling northern long eared bats and little brown bats in the greater Fundy ecosystem, New Brunswick. *Journal of Wildlife Management* 70: 5.
- Brunton, D.F. 1999. Natural environment inventory and interpretative concepts: Petrie Island study area, Cumberland, Ontario. 105 pp.
- Brunton, D.F. (2005) *City of Ottawa: Urban Natural Areas Environmental Evaluation Study – Appendix A – Vascular plants of the city of Ottawa with the identification of significant species*. Prepared for the Environmental Management Division Planning & Growth Management Department, City of Ottawa.
- City of Ottawa (2008). Official Plan – As adopted by Council – May 2008. x + 229pp
- City of Ottawa (2019). Significant Woodlands: Guidelines for Identification, Evaluation and Impact Assessment. 65pp.
- COSEWIC. (2003). COSEWIC assessment and status report on the Butternut *Juglans cinerea* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vii + 32 pp.

COSEWIC. (2005). COSEWIC assessment and update status report on the Blanding's Turtle *Emydoidea blandingii* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. viii + 40 pp.

COSEWIC. (2007). COSEWIC assessment and update status report on the Chimney Swift *Chaetura pelagica* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vii + 49 pp.

COSEWIC. (2009a). COSEWIC assessment and update status report on the Least Bittern *Ixobrychus exilis* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vi + 36 pp.

COSEWIC. (2010). COSEWIC assessment and status report on the Bobolink *Dolichonyx oryzivorus* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vi + 42 pp.

COSEWIC. (2013). COSEWIC assessment and status report on the Bank Swallow *Riparia riparia* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. ix + 48 pp.

Environment Canada. (2020). National Climate Data and Information Archive – OTTAWA INTL A. Accessed Online January 28, 2021 from: <http://climate.weatheroffice.gc.ca>.

Humphrey, C. (2017). Recovery Strategy for the Eastern Small-footed Myotis *Myotis leibii* in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Resources and Forestry, Peterborough, Ontario. vii + 76 pp.


Menzel, M, S. Owen, W. Edwards, P. Wood, B. Chapman & Miller, K. (2002). Roost tree selection by northern long-eared bat (*Myotis septentrionalis*) maternity colonies in an industrial forest of the central Appalachian Mountains. *Forest Ecology and Management* 155:107-114.

Muncaster Environmental Planning Inc. (2004). Wetland Impact Study – North Service Road Properties, Cumberland Ward, City of Ottawa. Report prepared for D.G. Belfie Planning & Development Consulting Ltd. 13 p.

Natural Resources Canada [EODMS \(nrcan-rncan.gc.ca\)](https://www.nrcan.gc.ca/eodms)


- OMNR. (2001). Ecological land Classification for Southern Ontario: Training Manual SCSS TM 01, March 2001.
- OMNR. (2010). Natural Heritage Reference Manual for Natural Heritage Policies of the Provincial Policy Statement, 2005. Ontario Ministry of Natural Resources. Second Edition: xi + 233 pp
- OMNR. (2013a). Ontario Wetland Evaluation System 3<sup>rd</sup>. Edition Version 3.3. viii + 284pp.
- OMNRF. (2013b). General Habitat Description for the Eastern Whip-poor-will (*Sturnella magna*). [Eastern Meadowlark General Habitat Description | Ontario.ca](#)
- OMNRF. (2014a). Land Information Ontario.
- OMNRF. (2015). Significant Wildlife Habitat Criteria Schedules for Ecoregions 6E. Ontario Ministry of Natural Resources and Forestry, Regional Operations Division, Peterborough. i + 38 pp.
- OMNRF. (2018). Bobolink General Habitat Description. Accessed Online January 23, 2019 from: <https://www.ontario.ca/page/bobolink-general-habitat-description>
- OMNRF. (2018). General Habitat Description for the Eastern Meadowlark (*Sturnella magna*). Accessed Online January 23, 2019 from: [http://files.ontario.ca/environment-and-energy/species-at-risk/mnr\\_sar\\_ghd\\_est\\_mdwrk\\_en.pdf](http://files.ontario.ca/environment-and-energy/species-at-risk/mnr_sar_ghd_est_mdwrk_en.pdf)
- Ontario Provincial Policy Statement. (2020).
- Peterson, R.T. (1980). *A field guide to the birds: A completely new guide to all the birds of eastern and central North America*. Houghton Mifflin Company, Boston.
- Sandilands, A. (2005). *Birds of Ontario Habitat Requirements, Limiting Factors and Status. Nonpasserines: waterfowl through cranes*. UBC Press Vancouver, BC. 260-263pp.
- Yates, M. D., and R. M. Muzika. 2006. Effects of Forest Structure and Fragmentation on Site Occupancy of Bat Species in Missouri Ozark Forests. *Journal of Wildlife Management*. Dec 2006: Vol 70, Issues 5, pp. 1238-1248


Appendix A : Background Review Mapping











Appendix B: Review of Potential for Significant Wildlife Habitat

Significant Wildlife Habitat	Candidate SWH ELC Codes	Candidate SWH Additional Criteria Summary	Confirmed SWM		Comments
			In Site	In Adjacent Lands	
<b>Seasonal Concentration Areas of Animals</b>					
Waterfowl stopover and staging areas (terrestrial)	Certain cultural meadow or thicket <u>Plus</u> evidence of annual spring flooding	Fields flooded from mid-March to May	No fields present with annual spring flooding.		Not Present; Not discussed further
Waterfowl stopover and staging areas (aquatic)	Specific aquatic habitat types (marsh, swamps)	Ponds, marshes, lakes, bays, coastal inlets and watercourses used for migration. Stormwater and sewage management facilities are not included.	No aquatic marshes, swamps etc.		Not Present; Not discussed further
Shorebird migratory stopover area	Beach/Bar Sand Dunes Meadow marsh	Shorelines used in May -t mid-June and early July to October. Stormwater and sewage management facilities are not included.	No shorelines, beaches, bars, dunes, or meadow marshes		Not Present; Not discussed further
Raptor wintering area	Requires combination of forest (deciduous, mixed or coniferous) and upland (cultural meadow, cultural thickets, cultural savannahs or cultural woodlands)	Combination of habitats must >20 ha and the field portion must be wind swept with little accumulation of snow. Where site is for eagles, open water and large trees and snags must be available.	No suitable forests are present		Not discussed further
Bat hibernacula	Crevice and caves	Active mines are not to be included. Buildings are not included.	No crevices or caves present		Not Present; Not discussed further

Significant Wildlife Habitat	Candidate SWH ELC Codes	Candidate SWH Additional Criteria Summary	Confirmed SWM		Comments
			In Site	In Adjacent Lands	
Bat maternity colonies	Deciduous, or mixed forests Deciduous or mixed Swamps (>5m tall)	>10/ha large diameter (>25 cm diameter at breast height) Snag trees in the decay classes 1-3 are preferred.	No forests are present in or within 15 m of the area to be disturbed. The edge trees along the ravine are smaller diameter.		Not Present; Not discussed further
Turtle wintering areas	Swamps, marshes, open water, shallow water, open fen or open bog	Water that is deep enough not to freeze solid with soft bottoms.  Must be permanent waterbody (or wetlands with adequate dissolved oxygen)	No ponds or other aquatic habitat present		Not Present; Not discussed further
Reptile hibernaculum	Any habitat except very wetlands Talus, rock barren, cave and alvar	For snakes – needs to be below frost lines.	Site consists of the disturbed lands.		Not Present; Not discussed further
Colonially – Nesting bird breeding habitat (Bank and Cliff Swallow)	Exposed sandy slopes of banks or piles. Cliff faces or structures (bridges, silos etc....)	Does not include licensed aggregate areas.  Does not include man-made structures or recently (within 2 years) disturbed soil	No exposed banks or cliffs present.		Not Present; Not discussed further
Colonially – Nesting bird breeding habitat (Trees/Shrubs)	Swamps – deciduous or mixed (trees >5m) Treed fen	Typically requires tall trees as nests are usually 11-15m from ground but shrubs and emergent vegetation could be used.	No swamps or treed fens present.		Not Present; Not discussed further

Significant Wildlife Habitat	Candidate SWH		Confirmed SWM		Comments
	ELC Codes	Additional Criteria Summary	In Site	In Adjacent Lands	
Colonially – Nesting bird breeding habitat (Ground)	Any rocky island or peninsula on lake or large river. For Brewer’s Blackbird – near watercourses in open fields, pastures		No rocky islands, or peninsulas were present. No suitable habitat for Brewer’s Blackbird were present.		Not Present; Not discussed further
Migratory butterfly stopover area	Not applicable to Ottawa Area – must be within 5 km of Lake Ontario				
Landbird migratory stopover area					
Deer yarding areas	Mixed or coniferous forests or swamps (>5m tall trees)  Can include plantations, cultural thickets, or dry-fresh poplar-white birch deciduous forest	These are mapped by OMNRF	None mapped by OMNRF for this area		Not Present; Not discussed further
Deer winter congregation area	All forest and wetland habitats and small conifer plantations	These are mapped by OMNRF (typically >100ha in size)			Not Present; Not discussed further
<b>Rare Vegetation Communities or Specialized Habitat for Wildlife</b>					
Cliffs and talus slopes	Near vertical face that is >3m in height (cliff or talus)	Typically in Niagara Escarpment	Cliffs and talus slope habitat were not present		Not Present; Not discussed further
Sand barren	Sand barrens various types but tree cover is always ≤ 60%	Must be >0.5ha	Sand barrens not present		Not Present; Not discussed further

Significant Wildlife Habitat	Candidate SWH ELC Codes	Candidate SWH Additional Criteria Summary	Confirmed SWM		Comments
			In Site	In Adjacent Lands	
Alvar	Alvar, Coniferous forest, cultural meadow, cultural savannah, cultural thickets and cultural woodlands	Must have at least 4 indicator species with substantial cover (must not have large amounts of exotic or introduced species)  Must be >0.5ha	Alvar habitat is typically flat and mostly unfractured calcareous bedrock. Not present		Not Present; Not discussed further
Old growth forest	Any forest or treed (>5 m) swamp	Must be at least 30 ha with at least 10 ha of interior habitat (edge considered 100 m)  Have specific characteristics (snags, mosaic of gaps, multi-layered canopy)	No forest present		Not Present; Not discussed further
Savannah	Tallgrass prairie savannah and cultural savannah	Must have indicator species	No savannah present		Not Present; Not discussed further
Tallgrass prairie	Tallgrass prairie (open prairie - <25% tree cover)	No minimum size	No tallgrass prairie were present. All area is manicured for multi-use pathway		Not Present; Not discussed further
Other rare vegetation communities	Provincially rare S1-S3 communities as described in Appendix M of the SWHTG		None of the communities listed for the Ottawa-Carleton Area in Appendix M were present.		Not Present; Not discussed further
<b>Specialised Habitat for Wildlife</b>					
Waterfowl nesting area	Shallow marsh, meadow marsh, thicket swamp or deciduous (treed >5 m tall) swamps	Wetland must be 0.5 ha or consist of up to 3 smaller wetlands within 120 m of each other if known nesting is occurring.	No marsh or swamp habitat present that was 0.5 ha. Base of ravine is less than 0.2 ha		Not Present; Not discussed further
Bald Eagle and Osprey nesting,	Any forest or swamp (trees >5m) type of habitat that is	Nests on man-made structures are not included.	No large trees and no Forests or treed Swamps were present on-site.		Not Present; Not discussed further

Significant Wildlife Habitat	Candidate SWH ELC Codes	Additional Criteria Summary	Confirmed SWM		Comments
			In Site	In Adjacent Lands	
foraging and perching habitat	immediately next to rivers, lakes, ponds or wetlands				
Woodland raptor nesting habitat	Any forest habitat or treed swamp (>5m tall) or coniferous plantation	Stand must be > 30 ha with >10 ha of interior habitat (edge is 200 m)	Does not meet the minimum requirements.		Not Present; Not discussed further
Turtle nesting areas	Shallow marsh, shallow water, open bog	Close to water but away from roads.  It must provide sand and gravel that turtles can dig through and be in open sunny areas.  Areas on the sides of municipal or provincial roads are not included.	Not present, pathway is paved.		Not Present; Not discussed further
Seeps and springs	Any forested community could have a seep/spring	Forest area with <25% meadow/pasture in the headwaters of a stream.		None present	Not Present; Not discussed further
Amphibian breeding habitat (woodland)	Any forest or treed swamp (>5m tall trees)	Wetland, pond or vernal pool must be > 500 m <sup>2</sup>  Those with water until mid-July (during most years) are better candidates		No wetlands or forests present	Not Present; Not discussed further
Amphibian breeding habitat (wetlands)	Swamps, marsh, fen, bog, open water or shallow water	Unless it's a larger wetland, must be >120 m from woodlands  Must be > 500 m <sup>2</sup>		No wetlands present	Not Present; Not discussed further

Significant Wildlife Habitat	Candidate SWH		Confirmed SWM		Comments
	ELC Codes	Additional Criteria Summary	In Site	In Adjacent Lands	
Woodland area-sensitive bird breeding habitat	Any forest or treed swamp (>5 m tall)	Interior habitat (200m edge used) in mature (>60 years) large (>30 ha) stand	No forest interior habitat present		Not Present; Not discussed further
<b>Habitat for Species of Conservation Concern (not including Endangered or Threatened Species)</b>					
Marsh bird breeding habitat	Meadow marsh, shallow water, fen or open bog		No marshes, shallow water or bogs present		Not Present; Not discussed further
Open country bird breeding habitat	Cultural meadows	Must be large grasslands (>30 ha)  Agricultural class 1 and 2 are not included  Agricultural lands planted in row crop or intensive hay, or pastures (within past 5 years) not included.	No grassland habitat present. Cultural meadow is disturbed, and vegetation is mostly broadleaf.		Not Present; Not discussed further
Shrub/early successional bird breeding habitat	Cultural thickets or woodlands	Must be > 10 ha  Agricultural class 1 and 2 are not included  Agricultural lands planted in row crop or intensive hay, or pastures (within past 5 years) not included	No thickets or woodlands are present		Not Present; Not discussed further
Terrestrial crayfish				Not present in Ottawa Area	
Special concern and rare wildlife species	All special concern or species ranked as S1-S3, SH (plants or animals)	Habitat depends on the species. Of those listed in <b>Error! Reference</b>		None.	Not Present; Not discussed further



Significant Wildlife Habitat	ELC Codes	Candidate SWH Additional Criteria Summary	Confirmed SWM In Site In Adjacent Lands	Comments
<b>source not found.</b> there is a potential for Snapping Turtle and Monarch.				
<b>Animal Movement Corridors</b>				
Amphibian movement corridor	Any habitat but amphibian breeding <u>wetland</u> habitat must be identified		Corridors need link habitats; upstream of this ravine is fully developed	Not Present; Not discussed further
Deer movement corridor	All forests but project must be in Stratum II Deer Wintering Area and Deer Wintering Habitat must be confirmed.		Not applicable – no Deer Wintering Areas or Habitat identified by OMNRF for area.	Not Present; Not discussed further

Appendix C: SAR Hand-Out

The following table provides photographs and general descriptions of potential species at risk that may occur within the project area and information on what actions to take should any of these species be observed.

Endangered and Threatened species are protected and cannot be harmed, harassed or killed and in some cases their habitats are also protected. These individuals will only be handled by qualified person and only if the individual is in imminent threat of harm. An authorization under the ESA 2007 would be required to handle individuals that are not in imminent threat of harm.

For all Endangered or Threatened species found on-site any activity which may cause harm to the individual will be stopped and the site supervisor will be contact immediately for further instructions.

### Chimney Swift



Photo: Mark Peck

[http://www.rom.on.ca/ontario/risk.php?doc\\_type=fact&lang=&id=322](http://www.rom.on.ca/ontario/risk.php?doc_type=fact&lang=&id=322)

### Description

A dark coloured bird with a light throat that has a cigar-shaped, cylindrical body with a short tail and long narrow wings.

**THREATENED**

### Action

Following is for both bird species:

- Stop any activity that may cause harm to these species and contact supervisor staff (see above)
- Individuals should only be encouraged to move if it is in immediate harm's way.

### Barn Swallow



Photo: Royal Ontario Museum website  
<http://www.rom.on.ca/ontario/fieldguides.php>

- Swallow with a long tail which is deeply forked in adult males
- 
- An orange front (no white on the forehead)
- Narrow pointed wings
- Juveniles have a white band across the top of the tail.

**THREATENED**

## Blanding's Turtle



Photo: Royal Ontario Museum website  
<http://www.rom.on.ca/ontario/risk.php>



Bernie Muncaster

### Description and Status

- Medium sized turtle (12.5-28 cm)
- Bright yellow on chin and throat.
- Shell is dark and can have light coloured spots or lines. The spots fade with age.
- The shell is domed.

### THREATENED

### Biology

- Lives in waterbodies – most often in areas with aquatic vegetation. But because this turtle moves very large distances through all kinds of habitats it can be encountered almost anywhere.
- Hibernates in water that is deep enough that it doesn't freeze to the bottom.
- It travels to get to or from the hibernation area, to find a mate or to lay its eggs. The hatchlings migrate towards water.
- The females do not tend to the eggs.
- They leave the hibernation sites in early spring (late April to mid-May).
- Can nest in gravel along road shoulders. Nests during late May to early June. Usually overnight or in early morning.
- Hatchlings leave the nest in the fall

#### *Types of Encounters:*

- Blanding's might travel through the area.
- They could nest in the road shoulder or on spoil piles.