

# **652 Flagstaff Drive – Environmental Impact Study**

**Ottawa, Ontario.**

*Palmer Project #*  
2301501

*Prepared For*  
Nukk Properties

December 18, 2023

December 18, 2023

Ram Balakrishnan  
Nukk Properties  
3812 Greenbank Road  
Ottawa, ON K2J 7B8

Dear Ram Balakrishnan:

**Re: 652 Flagstaff Drive – Environmental Impact Study**  
**Project #: 2301501**

---

Palmer is pleased to submit the attached Environmental Impact Study (EIS) for the proposed development at 652 Flagstaff Drive in the northeast corner of Borrisokane Road and Flagstaff Drive in Ottawa, Ontario. The Subject Property, approximately 0.4 ha, is currently a vacant lot occupied by regenerating cultural meadow vegetation and anthropogenic spaces.

The findings of our study are the result of a background review, field investigation, and an analysis of data using the current scientific understanding of the ecology of the area, as well as the current natural heritage policy requirements. We have identified the environmental sensitivities, constraints, and development opportunities of the Subject Property.

Based on the findings and recommendations of this study, it is our professional opinion that with the implementation of the mitigation measures provided in this report, that the proposed development plan adheres to the existing environmental framework and will not have negative impacts on natural features or their ecological functions.

Please let us know if you have question or comments on this submission.

Yours truly,

**Palmer™**



---

Austin Adams, M.Sc., EP  
Senior Ecologist

# Table of Contents

<b>1.</b>	<b>Introduction .....</b>	<b>1</b>
<b>2.</b>	<b>Environmental Policy .....</b>	<b>3</b>
2.1	Migratory Birds Convention Act (1994).....	3
2.2	Endangered Species Act (2007) .....	3
2.3	Provincial Policy Statement (2020) .....	3
2.4	City of Ottawa Official Plan (2021) .....	5
2.5	Rideau Valley Conservation Authority (O/Reg. 174/06).....	8
<b>3.</b>	<b>Background Review and Methodology .....</b>	<b>10</b>
3.1	Background Review .....	10
3.2	Ecological Surveys.....	10
3.2.1	Vegetation Communities and Flora.....	10
3.2.2	Wildlife, Species at Risk Habitat and Significant Wildlife Habitat Assessment.....	11
<b>4.</b>	<b>Existing Conditions.....</b>	<b>13</b>
4.1	Environmental Designation .....	13
4.2	Ecological Land Classification.....	13
4.3	Flora .....	16
4.4	Wildlife .....	16
4.5	Drainage Feature .....	16
<b>5.</b>	<b>Assessment of Significance.....</b>	<b>18</b>
5.1	Species At Risk.....	18
5.2	Significant Wildlife Habitat.....	18
<b>6.</b>	<b>Proposed Development .....</b>	<b>20</b>
<b>7.</b>	<b>Impact Assessment and Mitigation Measures .....</b>	<b>22</b>
7.1	Woodland and Wetland.....	22
7.2	Drainage Feature .....	22
7.3	Stormwater Management.....	24
<b>8.</b>	<b>Policy Conformity.....</b>	<b>25</b>
<b>9.</b>	<b>Conclusions .....</b>	<b>26</b>
<b>10.</b>	<b>Certification .....</b>	<b>27</b>
<b>11.</b>	<b>References .....</b>	<b>28</b>

## List of Figures

---

Figure 1. Site Location.....	2
Figure 2. Existing Conditions.....	12
Figure 3. Proposed Development.....	21

## List of Tables

---

Table 1. Policy Conformity .....	25
----------------------------------	----

## List of Appendices

---

Appendix A. Species at Risk Screening	
Appendix B. Plant List	
Appendix C. Significant Wildlife Habitat Screening	

# 1. Introduction




Palmer has been retained by Nukk Properties to complete an Environmental Impact Study (EIS) for the property identified as 652 Flagstaff Drive, located in the northeast corner of Borrisokane Road and Flagstaff Drive, City of Ottawa (the Subject Property, **Figure 1**). The proponent wishes to develop the Subject Property for 14,800 sq. ft. retail/restaurant space in two 1-storey commercial buildings, with 68 parking stalls.

The Subject Property is currently a vacant lot occupied by regenerating cultural meadow vegetation and anthropogenic spaces. A watercourse is located east of the Subject Property and flows north towards Jock River.



This EIS relies on background information and a single site visit by a Palmer ecologist to investigate any potential impacts or environmental policy implications associated with this proposed development. The property occurs within the planning area for the City of Ottawa, as well as the Rideau Valley Conservation Authority (RVCA) regulated area.



LEGEND

-  Watercourse <sup>1</sup>
-  Wetland - Unevaluated as per OWES <sup>2</sup>
-  Subject Property

1 - Ontario Hydro Network (OHN)  
2 - Land Information Ontario (LIO)



North American Datum 1983  
Universal Transverse Mercator Projection Zone 17

Scale: 1:2,000  
Page Size: Tabloid (11 x 17 inches)

Drawn: SM  
Checked: HB  
Date: Aug 23, 2023



Source Notes:  
Base imagery (2021) provided by Ontario gis rest services.  
Contains information licensed under the Open Government Licence - Ontario.

CLIENT	Nukk Properties
PROJECT	Flagstaff Borrisoakane
TITLE	<b>Site Location</b>
REF. NO.	2301501-1-1
<b>Figure 1</b>	

## 2. Environmental Policy

The environmental policies applicable to the Subject Property have been reviewed with specific relevant policies summarized in the following sections. The environmental policies of the Province, City of Ottawa Official Plan (OP), RVCA development policies, the provincial *Endangered Species Act, 2007* (ESA), and the *Migratory Birds Convention Act, 1994* have all been considered.

### 2.1 Migratory Birds Convention Act (1994)

The *Migratory Birds Convention Act, 1994* (MBCA) and *Migratory Birds Regulations, 2022* (MBR) protect most species of migratory birds and their nests and eggs anywhere they are found in Canada (Government of Canada, 1994). General prohibitions under the MBCA and MBR protect migratory birds, their nests and eggs and prohibit the deposit of harmful substances in waters / areas frequented by them. The MBR includes an additional prohibition against incidental take, which is the inadvertent harming or destruction of birds, nests, or eggs.

Compliance with the MBCA and MBR is best achieved through a due diligence approach, which identifies potential risk, based on a site-specific analysis in consideration of the Avoidance Guidelines and Best Management Practices information on the Environment Canada website (Government of Canada, 2018).

### 2.2 Endangered Species Act (2007)

Species designated as Endangered or Threatened by the Committee on the Status of Species at Risk in Ontario (COSSARO) are listed as Species at Risk in Ontario (SARO). These species at risk (SAR) and their habitats (e.g., areas essential for breeding, rearing, feeding, hibernation, and migration) are afforded legal protection under the *Endangered Species Act, 2007* (ESA) (Government of Ontario, 2007). This Act is administered by the Ministry of Environment, Conservation and Parks (MECP).

The protection provisions for species and their habitat within the ESA apply only to those species listed as Endangered or Threatened on the SARO list, being Ontario Regulation 230/08 of the ESA. Species listed as Special Concern may be afforded protection through policy instruments respecting significant wildlife habitat (e.g., the Provincial Policy Statement (PPS)) as defined by the Province or other relevant authority, or other protections contained in municipal Official Plan policies.

### 2.3 Provincial Policy Statement (2020)

The *Provincial Policy Statement* (PPS) provides direction to regional and local municipalities regarding planning policies for the protection and management of natural heritage features and resources (Ontario Ministry of Municipal Affairs and Housing, 2020). The PPS defines eight types of Natural Heritage Features (NHF) and adjacent areas and provides planning policies for each. Of these NHF, development is not permitted in:

- Significant Coastal Wetlands;
- Significant Wetlands in Ecoregions 5E, 6E and 7E;
- Fish Habitat, except in accordance with provincial and federal requirements; or

- Habitat of species designated as Endangered and Threatened, except in accordance with provincial and federal requirements.

Additionally, unless it can be demonstrated through an Environmental Impact Study (EIS) that there will be no negative impacts on the natural features or their ecological functions, development and site alteration are also not permitted in:

- Significant Wetlands in the Canadian Shield north of Ecoregions 5E, 6E and 7E;
- Significant Woodlands in Ecoregions 6E and 7E (excluding islands in Lake Huron and the St. Mary's River);
- Significant Valleylands in Ecoregions 6E and 7E (excluding islands in Lake Huron and the St. Mary's River);
- Significant Wildlife Habitat;
- Significant Areas of Natural and Scientific Interest (ANSI);
- Other Coastal Wetlands in Ecoregions 5E, 6E and 7E; and
- Lands defined as *Adjacent Lands* to all the above natural heritage features.

The Subject Property occurs in Ecoregion 6E, as shown below in **Map A**, and does not contain any of the abovementioned features. Woodland and wetland communities are noted as occurring to the south.



**Map A: Natural Heritage Information Centre (NHIC) mapping depicting a watercourse (blue line) to the south and east of the Subject Property (approximate boundaries in red). Unevaluated wetlands (blue patterned layer) and woodlands (green layer) are noted south of the Subject Property.**



## 2.4 City of Ottawa Official Plan (2021)

The purpose of the City of Ottawa Official Plan (2021) is to “manage this growth in ways that reinforce the qualities of the city most valued by its residents: its distinctly liveable communities, its green and open character, and its unique characteristics that distinguish Ottawa from all other places”.

According to Section 4.8.1 of the City’s OP:

1) *The Natural Heritage System consists of core natural areas and natural linkage areas. Natural Heritage Features occur both inside and outside the Natural Heritage System. The Natural Heritage System and the features within it are subject to a higher standard of protection than features outside the Natural Heritage System.*

3) *The City recognizes the following natural heritage features, as defined in Ottawa’s Environmental Impact Study Guidelines:*


- a) *Significant wetlands;*
- b) *Habitat for endangered and threatened species;*
- c) *Significant woodlands;*
- d) *Significant valleylands*
- e) *Significant wildlife habitat;*
- f) *Areas of Natural and Scientific Interest;*
- g) *Urban Natural Features;*
- h) *Natural Environment Areas;*
- i) *Natural linkage features and corridors;*
- j) *Groundwater features;*
- k) *Surface water features, including fish habitat; and*
- l) *Landform features.*

4) *The natural heritage overlay policies apply to all features in Policy 3) regardless of whether they appear on Schedules to the Official Plan.*

As per the City of Ottawa’s *Environmental Impact Statement Guidelines* (2015) definitions, none of these Natural Heritage Features are present on the Subject Property. This was confirmed through the June 2023 field investigation and the review of aerial imagery and OP mapping. Natural Heritage Features; however, do potentially occur on Adjacent Lands and are discussed later within this report.

Schedule C11-B of the OP mapping (**Map B**) indicated that the Subject Property is within an Urban Area and does not contain Natural Heritage System Core Areas, Natural Heritage System Linkage Areas, or Natural Heritage Features. As such, the Subject Property is not subject to Natural Heritage System Core Areas, Natural Heritage System Linkage Areas, or Natural Heritage Features policies. A watercourse is noted immediately west of the Subject Property, and two additional watercourses are documented to the south and the east.

Schedule C12 of the OP mapping (**Map C**) indicated that the Subject Property, which is within Ottawa’s Urban Area, does not overlap with the Urban Greenspace network. Additionally, the Subject Property does not contain any Greenbelt Natural Areas or Greenbelt Natural Linkage Areas. This is also reflected in Schedule B4 – Greenbelt Transect of the City’s OP. As such, the Subject Property is not subject to Urban Greenspace, Greenbelt Natural Areas, or Greenbelt Natural Linkage Areas policies. An Urban Natural Feature (UNF) is noted in lands south of the Subject Property. An Open Space is also noted on lands immediately southeast of the Subject Property.



*Map B: Schedule C11-B Natural Heritage System (South) - Subject Property (approximate location denoted by red arrow) located within the Urban Area land designation (yellow layer) with watercourses to the west, east and south of the site, and an Urban Natural Feature / Natural Heritage Feature to the south (green layer outlined in purple).*




*Map C: Schedule C12 Urban Greenspace - Subject Property (boundaries in red) located within the Urban Area land designation (white layer) with a UNF to the south (dark green layer) and an Open Space (light green layer) immediately to the southeast.*

As per Section 10.1.1 of the City's OP:

*Lands in the flood plain are subject to regulations under Section 28 of the Conservation Authorities Act, Official Plan policies and zoning provisions in the Zoning By-law. Official Plan policies for flood plains also apply to areas where flood plains exist but are not mapped by the conservation authorities or identified as flood plain in the Zoning By-law. Flood plain boundaries are determined in consultation with the Mississippi Valley, Rideau Valley and South Nation Conservation Authorities, with adherence to relevant policies and guidelines. Schedule C15 – Environmental Constraints provides the general location of lands affected by the flood plain, however this schedule does not show all lands affected by the flood plain and should not be used to determine if a lot is affected by the flood plain. For detailed information about lands affected by the flood plain, refer to the Flood Plain Overlay in the Zoning By-law. For areas where no flood plain mapping is available, development proponents may be required to undertake studies as part of the development review and approvals process to delineate the extent of the flood plain.*

According to Schedule C15 of the OP mapping (**Map D**) the Subject Property is located within a flood plain. This is further confirmed through RVCA mapping (see **Map E** in Section 2.5 below), whereby regulated lands transect the Subject Property. Thus, the Subject Property is subject to floodplain regulations under Section 28 of the Conservation Authorities Act, Official Plan policies, and zoning provisions in the Zoning By-law.



**Map D: Schedule C15 Environmental Constraints – Subject Property located within Flood Plain (teal layer) and Organic Soil (orange layer) to the south of the Subject Property (approximate location denoted by red arrow).**

## 2.5 Rideau Valley Conservation Authority (O/Reg. 174/06)

The Subject Property occurs within the jurisdiction of RVCA, regulated under O/Reg. 174/06 (Rideau Valley Conservation Authority, 2013).

Regarding development prohibition:

*2. (1) Subject to section 3, no person shall undertake development or permit another person to undertake development in or on the areas within the jurisdiction of the Authority that are,*

*(a) adjacent or close to the shoreline of inland lakes that may be affected by flooding, erosion or dynamic beaches, including the area from the furthest offshore extent of the Authority's boundary to the furthest landward extent of the aggregate of the following distances:*

*(i) the 100 Year flood level,*

*(ii) the predicted long term stable slope projected from the existing stable toe of the slope or from the predicted location of the toe of the slope as that location may have shifted as a result of shoreline erosion over a 100-year period, and*

*(iii) an allowance of 15 metres inland;*

*(b) river or stream valleys that have depressional features associated with a river or stream, whether or not they contain a watercourse, the limits of which are determined in accordance with the following rules:*

*(i) where the river or stream valley is apparent and has stable slopes, the valley extends from the stable top of bank, plus 15 metres, to a similar point on the opposite side,*

*(ii) where the river or stream valley is apparent and has unstable slopes, the valley extends from the predicted long term stable slope projected from the existing stable slope or, if the toe of the slope is unstable, from the predicted location of the toe of the slope as a result of stream erosion over a projected 100-year period, plus 15 metres, to a similar point on the opposite side,*

*(iii) where the river or stream valley is not apparent, the valley extends the greater of,*

*(A) the distance from a point outside the edge of the maximum extent of the flood plain under the applicable flood event standard, plus 15 metres, to a similar point on the opposite side, and*

*(B) the distance from the predicted meander belt of a watercourse, expanded as required to convey the flood flows under the applicable flood event standard, plus 15 metres, to a similar point on the opposite side;*

*(c) hazardous lands;*

*(d) wetlands; or*

*(e) other areas where development could interfere with the hydrologic function of a wetland, including areas within 120 metres of all provincially significant wetlands and wetlands greater than 2 hectares in size, and areas within 30 metres of wetlands less than 2 hectares in size. O. Reg. 174/06, s. 2 (1); O. Reg. 78/13, s. 1 (1, 2).*

As shown below on **(Map E)**, the Subject Property does contain regulated areas, identified hazard areas, or natural heritage features as shown in online web mapping by the RVCA. Under O. Reg. 174/06, a permit would be required from the RVCA prior to development within these regulated lands.



**Map E: RVCA Mapping showing RVCA Regulated Areas (green hatching layer) and watercourse (blue line), and wetland (green polygon layer) to the southeast of the Subject Property (approximate boundaries in red).**

## 3. Background Review and Methodology

### 3.1 Background Review

Palmer has reviewed relevant background material to provide a focus on field investigations and ensure compliance with applicable regulations and policy. Background information collection is guided by the *Natural Heritage Information Request Guide* (Ministry of Natural Resources and Forestry, 2018). Current direction from the Ministry of Natural Resources and Forestry (MNRF) and Ministry of Environment, Conservation and Parks (MECP) is to gather natural heritage information and species occurrence records from available sources; the Natural Heritage Information Centre (NHIC) Make-a-Map application being the main source of information and records from the Ministry itself (Ministry of Natural Resources and Forestry, 2023). Information gathered is recommended to be balanced and supplemented by a professional ecological review of potential habitats and characteristics of a project site.

Background review included the collection and review of relevant mapping and reports, including regulations and policies, Official Plans, and the NHIC Make-a-Map application for species occurrences and designated area mapping. In addition to these sources, the following data sources were reviewed for the project:

- **Land Information Ontario (LIO):** Certain data types including aquatic resource area (ARA) information is available through these publicly available data layers (Government of Ontario, 2023).
- **Conservation Authority:** The RVCA collects and maintains natural heritage mapping and data, and publish reports, that all provide regional and often site-specific ecological context.
- **Ontario Breeding Birds Atlas (OBBA):** For breeding bird records in the general vicinity (Bird Studies Canada, 2023).
- **Ontario Reptile and Amphibian Atlas (ORAA):** Provides range maps and other information regarding reptile and amphibian species observed in Ontario (Ontario Nature, 2023).

Following the *Information Request Guide* (MNRF, 2018), MECP advice and direction should be solicited once Species at Risk (SAR) interactions or potential interactions are identified via field investigations and analysis. .

### 3.2 Ecological Surveys

An ecological field survey was conducted on June 5, 2023. The weather conditions at the time of Palmer's survey included overcast skies with a temperature of 22°C. The ecological investigation included in-field data collection for vegetation communities, to the extent possible during the spring season, as well as the review of recent aerial imagery.

#### 3.2.1 Vegetation Communities and Flora

Vegetation communities were mapped and described following the Ecological Land Classification (ELC) System for Southern Ontario (Lee, et al., 1998). Vegetation community boundaries were delineated on field

maps through the interpretation of recent aerial photographs and refined in the field. Information collected during ELC includes dominant species cover, community structure, as well as level of disturbance, presence of indicator species, and other notable features. The vegetation community polygons determined through aerial photos were more precisely defined and identified to ELC Vegetation Type or Ecosite or other best-fitting description. Dominant species, level of disturbance, and features of interest were recorded for each community.

Botanical surveys were completed by traversing the Subject Property and recording species observed in each vegetation community. Provincial plant status was based on the *Provincially Rare Flora of Ontario* (Oldham & Brinker, 2009) and the NHIC species lists (Ministry of Natural Resources and Forestry, 2023). Searches for Butternut (*Juglans cinerea*), an *Endangered* tree, were completed during the ELC and botanical surveys.

### 3.2.2 Wildlife, Species at Risk Habitat and Significant Wildlife Habitat Assessment

A desktop study was completed using the *Ontario Breeding Bird Atlas* (Bird Studies Canada, 2023), the *Ontario Reptile and Amphibian Atlas* (Ontario Nature, 2023) to determine bird and herpetofauna species that might occur on the Subject Property, respectively. All incidental observations of wildlife were recorded by Palmer during the investigations. Incidental observations included direct sightings and indirect evidence such as nests, tracks, scat, and browse.

An assessment for Species at Risk (SAR) habitat and Significant Wildlife Habitat (SWH) has been completed for the Subject Property through a review of potential SAR and SWH values known to occur in the region. The potential presence of SAR and SWH was determined through background review and observations from the single site investigation.

Prior to field investigations, a background review was completed for potential SAR habitat opportunities. The NHIC database, the Ontario Breeding Bird Atlas (OBBA), and the Ontario Reptile and Amphibian Atlas (ORAA) were screened for SAR records. The Subject Property includes NHIC square 1107866, which indicated that Butternut (*Juglans cinerea*), and Bobolink (*Dolichonyx oryzivorus*) are known to occur within or adjacent to the Subject Property. The Subject Property was assessed for potential SAR habitat opportunities during the June 5 site visit.

This assessment was conducted by comparing habitat preferences of species deemed to have potential to occur against current site conditions. This SAR habitat assessment can be found in **Appendix A**, providing a detailed description of each species' habitat (including those deemed to not have potential habitat), as well as a discussion of habitat suitability within the Subject Property, and potential impacts and mitigation where applicable. Based on the rationale provided in **Appendix A**, no potential SAR habitat was identified as potentially present on the Subject Property.



LEGEND

Watercourse <sup>1</sup>



Ecological Land Classification (ELC)

**ELC LEGEND:**  
 ANTH: Anthropogenic  
 CUM: Cultural Meadow

Wetland - Unevaluated as per OWES <sup>2</sup>

Subject Property

1 - Ontario Hydro Network (OHN)  
 2 - Land Information Ontario (LIO)



North American Datum 1983  
 Universal Transverse Mercator Projection Zone 17

Scale: 1:500  
 Page Size: Tabloid (11 x 17 inches)

Drawn: SM  
 Checked: HB  
 Date: Aug 23, 2023



Source Notes:  
 Base imagery (2021) provided by Ontario gis rest services.  
 Contains information licensed under the Open Government  
 Licence - Ontario.

CLIENT	Nukk Properties
PROJECT	Flagstaff Borrisoakane
TITLE	<b>Existing Environmental Condition</b>
REF. NO.	2301501-2-1
	<b>Figure 2</b>



## 4. Existing Conditions

### 4.1 Environmental Designation

Based on the review of the MNR's Make-a-map application, the Subject Property has no provincial environmental designations (i.e., ANSI, Provincially Significant Wetlands) (Ministry of Natural Resources and Forestry, 2023) (**Map A**). To the south but removed from the Subject Property is a large, naturalized block, where NHIC mapping shows the presence of a woodland and an unevaluated wetland (**Map A**). Additionally, drainage/watercourse features were noted on the south and east side of the Subject Property, adjacent to the Subject Property boundaries (**Map F**). This image shows that this watercourse is a manmade channel but has been engineered to include naturalized elements such as meanders and widened segments.



*Map F: Image showing constructed channel with naturalized design adjacent to the Subject Property (Microsoft, 2023).*

### 4.2 Ecological Land Classification

The Subject Property occurs on the east side of Borrisokane Road in an area of anthropogenic influence, including a suburban development to the east and historic farmlands on all other surrounding lands. The Subject Property was historically used for agricultural purposes but has been left fallow for several years. Currently, the primary vegetation cover is regenerating meadow species. No trees were observed on the Subject Property at the time of Palmer's survey.

The Subject Property occurs approximately 500 m south of the Jock River. There are identified site drainages to the west and east of the Subject Property which flow north towards this river (**Photos 1 & 2**). The eastern drainage is constructed, with natural elements (**Map F**), whereas the west side is a roadside ditch system. The Jock River then ultimately conveys waters east to the Rideau River.

The two ELC communities identified within the Subject Property are shown on **Figure 2** and is further described as follows:

#### Anthropogenic (ANT)

A small area in the west end of the Subject Property includes an existing gravel flat. Palmer expects that this area has recently been utilized for access or parking purposes. A small area in the south along the property line has exposed soils with no vegetation present (**Photo 3**).

#### Cultural Upland Meadow (CUM)

The only vegetated areas within the Subject Property were Cultural Upland Meadows, which appeared to be historically cleared and maintained for agricultural use but has since been left to fallow (as mentioned above). Portions of the meadow appeared to be sparse (**Photos 4 & 5**) with exposed soil. The topography of this community is slightly raised from Flagstaff Road (Approx. 0.5 m). Then slopes downward (Approx. 1.5 m) at the northern portion of the subject Property. The majority of the Subject Property is cultural meadow (CUM ELC polygon), which is the result of regenerating species after the clearing of agricultural lands and contains sparse grassy areas and forb species such as Red Clover (*Trifolium pratense*), White Clover (*Trifolium repens*), Birds-foot Trefoil (*Lotus corniculatus*), and Aster species (*Symphotrichum* sp.). No persistent stalks were observed during the site visit (which would indicate a well-established grassland).



***Photos 1 & 2: Photo 1 - East side of Subject Property - Drainage Feature for Jock River on adjacent (east) property (looking southeast). Photo 2 - Drainage Feature for Jock River on adjacent property to the west (looking north).***



*Photo 3: South portion of Subject Property (facing east) – with no vegetation present.*



*Photo 4: Subject Property with sparse meadow forbs. Primary Cultural Meadow (CUM) through Subject Property.*



*Photo 5: Subject Property with woodlands to the south.*

### 4.3 Flora

From spring and summer surveys, 32 species of vascular plants were recorded within the Subject Property, including 8 (25%) native species, 21 (66%) which are non-native to Ontario, and three species that were identified to the genus only due to the limited representation of key characteristics (**Appendix B**). The recorded presence of this percentage of non-native species is indicative of farming and past disturbance in the Subject Property, and is typical of developed areas in Southern Ontario (Morton & Venn, 1984). Similarly, Oldham et al. (1995) indicate that in southern Ontario plant communities, non-native flora presence averages between 20 and 30%.

All species observed are considered provincially secure (S5) or non-native/exotic species (NNA) by NHIC (Ministry of Natural Resources and Forestry, 2023). All species are Common in the Ottawa area, with the exception of Common Wormwood (*Artemisia vulgaris*), which is Uncommon, but local to the Ottawa River; this species is considered a non-native/exotic species by NHIC (Brunton, 2005; Ministry of Natural Resources and Forestry, 2023)

### 4.4 Wildlife

Red-winged Blackbird (*Agelaius phoeniceus*) was observed within and adjacent to the Subject Property during the June 5, 2023 site visit. This is a common species in Ontario and is likely using the adjacent naturalized watercourse as habitat. Due to the lack of established vegetation on the Subject Property. No other wildlife sightings were observed during the site visit.

### 4.5 Drainage Feature

The drainage features identified on the south and east side of the Subject Property in NHIC were identified with no water present during Palmer's 2023 site visit. Although a roadside ditch is identified along the western edge of the Subject Property, the 2023 site investigation noted that this ditch was completely dry.

During Palmer's site investigation, the only identified drainage feature was noted on a property to the east, approximately 40 metres from the eastern boundary of the Subject Property. This drainage feature flowed directly into the Jock River to the north and presented a potential risk for erosion and sediment runoff. This drainage feature was observed as dry during the time of the site visit.

## 5. Assessment of Significance

The significance of the identified potential ecological features on the Subject Property are evaluated below for their significance and applicability to the policies detailed in Section 2. Other Natural Heritage Features (e.g., Significant Valleylands) are not present on the Subject Property.

### 5.1 Species At Risk

No Butternut were detected on the Subject Property during the site visit. It can be confirmed that Butternut trees do not currently exist on or immediately adjacent to the Subject Property.

Several species of SAR birds occur in the region, according to Ontario Breeding Bird Atlas results, as shown in **Appendix A**. Even though the Subject Property has a meadow community, it does not include extensive cover by grassy vegetation, and therefore is not suitable habitat for these SAR bird species. Potential suitable habitat may be present immediately north of the Subject Property. A more established meadow community is located north of the Subject Property with persistent stalks observed.

Several species of SAR reptile are identified by the Ontario Reptile and Amphibian Atlas (ORAA), as shown in **Appendix A**, as occurring within the region. These species, however, would not find habitat within the Subject Property due to a complete lack of upland micro-habitat features (i.e., rock or shrub cover, woody debris) and aquatic habitat. These species may occur within the adjacent eastern watercourse and Jock River to the north, which is the likely the source of the historical records.

### 5.2 Significant Wildlife Habitat

Significant Wildlife Habitat (SWH) can be difficult to appropriately determine at the site-specific level, as the assessment must incorporate information from a wide geographic area and consider other factors such as regional resource patterns and landscape effects. To help with site level assessments, the MNR has developed the *Significant Wildlife Habitat Criteria Schedules for Ecoregion 6E* (Ontario Ministry of Natural Resources and Forestry, 2015).

SWH is defined by the MNR in the Significant Wildlife Habitat Technical Guide (Ontario Ministry of Natural Resources, 2000) and Natural Heritage Reference Manual (Ontario Ministry of Natural Resources, 2010) and includes the following categories:

- Seasonal Concentration Areas of Animals;
- Rare Vegetation Communities or Specialized Habitats for Wildlife;
- Habitats of Species of Conservation Concern; and
- Animal Movement Corridors.


Criteria for the identification of these features are also provided in the *Significant Wildlife Habitat Criteria Schedules for Ecoregion 6E*. These criteria were used to provide an assessment for wildlife habitat within and adjacent to the Subject Property, focusing primarily on lands within and immediately adjacent to the proposed development footprint. The SWH assessment is detailed in **Appendix C**.

No SWH were identified as having potential to occur within the Subject Property; however, SWH values likely occur in the adjacent natural communities.

## 6. Proposed Development

As detailed on the Site Plan prepared by Architecture Urban Design (dated April 2023), the proposed development involves 14,800 sq. ft. retail/restaurant space in two 1-storey commercial buildings, with 68 parking stalls on the Subject Property, with an associated access road (**Figure 3**). Access to the Subject Property will be facilitated via the proposed access road along the east side of the Subject Property, which connects to the newly developed access road off Flagstaff Drive. The proposed design has been revised to allow for a 3 m setback on the east side, allowing for a buffer to the adjacent watercourse.





LEGEND

Watercourse <sup>1</sup>

3 m Landscape Buffer

ESC Fence

Wetland - Unevaluated as per OWES <sup>2</sup>



Ecological Land Classification (ELC)

Development Plan

Subject Property

**ELC LEGEND:**  
 ANTH: Anthropogenic  
 CUM: Cultural Meadow

1 - Ontario Hydro Network (OHN)  
 2 - Land Information Ontario (LIO)



North American Datum 1983  
 Universal Transverse Mercator Projection Zone 17

Scale: 1:500  
 Page Size: Tabloid (11 x 17 inches)

Drawn: SM  
 Checked: HB  
 Date: Aug 23, 2023

Source Notes:  
 Base imagery (2021) provided by Ontario gis rest services.  
 Contains information licensed under the Open Government  
 Licence - Ontario.



CLIENT	Nukk Properties
PROJECT	Flagstaff Borrisoakane
TITLE	<b>Proposed Development</b>
REF. NO.	2301501-3-3
	<b>Figure 3</b>



## 7. Impact Assessment and Mitigation Measures

The Subject Property does not contain any natural features within its boundaries that require removal or modification for the proposed development. Natural features, have been identified on nearby lands, including a wetland and woodland to the south of the Subject Property and a realigned drainage feature/watercourse conveying north to the Jock River to the east of the Subject Property. Indirect impacts to adjacent and off-site features are the most significant potential effects of the proposed development to be considered, as the onsite cultural meadow has been identified as supporting mainly non-native species and provides limited, low-quality wildlife opportunities.

### 7.1 Woodland and Wetland

To the south of the Subject Property, a woodland/wetland complex occurs. This UNF is approximately 200 m south of the proposed Subject Property's south boundary. Consequently, this UNF is outside of the lands considered "Adjacent Lands" by the PPS (not within 120 m of the Subject Property) and, as such, no impacts are anticipated on this woodland/wetland complex.

Furthermore, the topography of the landscape on and adjacent to the Subject Property, as shown in **Photos 1-5**, is generally flat with overland conveyance of erosion. Although this erosion should not impact the UNF, standard mitigation measures during construction, as discussed below, should be implemented to ensure that there are no impacts to this woodland/wetland feature. After construction, no impacts are anticipated on this UNF given the distance and the nature of Subject Property's topography.

### 7.2 Drainage Feature

The drainage feature to the east has been realigned, which was completed as part of an adjacent property's development application. The City of Ottawa conducted a site visit on January 16, 2023 to inspect the Subject Property and the adjacent site and to verify the status of the realigned watercourse. The City of Ottawa confirmed that, as per provision 3 of Section 69 of the zoning by-law, that no formal setback from the drainage feature will be required on the Subject Property. Additionally, the realigned drainage feature must also provide an ecological and hydrological function that is similar to that of the old drainage feature. Consequently, the realigned feature must allow for amphibian transit between the UNF and the Jock River, must prevent fish from obtaining access to the wetland swamp (i.e., the Cambrian Woods North), and must not alter water levels within the wetland swamp (e.g., the UNF). These factors, as well as Erosion and Sediment Controls (ESC) and the proposed development's operations, were considered when determining appropriate mitigation measures for the adjacent realigned watercourse. Measures to mitigate any adverse impacts have been provided below.

#### Erosion and Sediment Control (ESC)

- Detailed ESC measures are outlined in Civil Plan C101 submitted as part of the development application, and the general configuration also shown on **Figure 3**. ESC measures aim to prevent movement of sediment laden stormwater generated on-site during construction activities into adjacent drainage systems. Increased sediment loads into aquatic habitats can have adverse

impacts such as increased water temperatures, a reduction in oxygen availability that in turn pose stress on resident fish populations, and increased turbidity which can impact the ability for wildlife to forage.

### Light Pollution

- A variety of native trees and shrubs are proposed to be planted along the property boundary on the north and east sides. To mitigate impacts from light sources on the proposed development. Some species include Dwarf Bush-honeysuckle (*Diervilla lonicera*), Northern Oak (*Quercus rubra*), Common Hackberry (*Celtis occidentalis*) and Fragrant Sumac (*Rhus aromatica*).
- Along with a mix of deciduous trees for canopy coverage, Eastern White Cedar (*Thuja occidentalis*) and native shrubs such as Snowberry (*Symphoricarpos*), Chokeberry (*Prunus virginiana*), Buttonbush (*Cephalanthus occidentalis*) and Gray Dogwood (*Cornus racemosa*) are proposed to be planted along the eastern boundary of the Subject Property to mitigate impacts from light sources on the proposed development. These plantings will provide shading for the adjacent realigned watercourse. Planting specifications are outlined in the Landscaping Plan L-01 submitted as part of the development application and will incorporate species approved by the City of Ottawa (Rhuland and Associates Ltd., 2023).
- Where space/widths allow in Landscaping Plan L-01, to increase the protection of the adjacent watercourse, conifers including Eastern White Cedar, Eastern Hemlock (*Tsuga canadensis*), Eastern White Pine (*Pinus strobus*), and White Spruce (*Picea glauca*) will be planted around the garage enclosure. Matching with other landscape requirements and objectives, this planting location will not interfere with sight lines or spread into the parking area.
- The eastern limits of the property (buffer planting plan area – **Figure 3**) are also recommended to be seeded with an appropriate mixture of native perennials, with the objective of achieving a layer of *native, self-sustaining vegetation*.

### Air and Noise Pollution

- Signage will be posted on the Subject Property to encourage customers not to idle while on the premises.
- As above, native trees/shrubs/plants will be planted along the eastern and surrounding boundary of the Subject Property to mitigate impacts from vehicular traffic, including idling (carbon monoxide, dust) and noise. Planting specifications are outlined in the Landscaping Plan submitted as part of the development application and will incorporate species approved by the City of Ottawa (Rhuland and Associates Ltd., 2023).

### Snow Removal

- Snow will be collected and removed from the Subject Property to an off-site storage area. This will ensure that road salts, sediments and other contaminants built up in snows will not reach the adjacent watercourse. Signs will be posted along the edge of the eastern boundary to ensure protection from snow clearing, management, storage and removal.

### Garbage

- An enclosure will be present on the Subject Property to store garbage. Specifications are outlined in Site Plan SP01 that was submitted as part of the development application.

### Protocol for Wildlife Protection during Construction (City of Ottawa, 2015)

- Due to the lack of established vegetation and natural heritage features on the Subject Property, the impact to potential wildlife is low in the form of habitat removal during construction and post-construction impacts to local wildlife. Regardless, mitigation measures should be implemented during construction activities to minimize any potential impacts. These mitigation measures include having proper waste control measures (i.e., proper garbage receptacles) and proper drainage measures to avoid attracting wildlife to the area. Additionally, clearing of the meadow species should begin on the west side of the Subject Property and progress east towards the realigned drainage feature, giving any potential wildlife the opportunity to move towards the natural feature.

### Jock River Reach 1 Subwatershed Plan (City of Ottawa, 2006)

- Implementation of the abovementioned mitigation measures will result in no negative impacts to the realigned drainage feature east of the Subject Property, which will consequently result in no negative impacts to Reach 1 of the Jock River. These mitigation measures will allow for naturalization along the new drainage feature corridor.

## **7.3 Stormwater Management**

The Subject Property occurs in an area that has a low capacity to buffer for soil erosion, as the Subject Property and its immediate surroundings are primarily devoid of rooted vegetation. The use of ESC (as mentioned above) during construction will be required to limit any potential impacts to adjacent properties or natural heritage features (i.e., the realigned drainage feature).

The project design indicates that stormwater from the rooftops and parking lots will be collected into catch basin manholes and conveyed through underground storm sewers, stored in underground storage chambers (as needed for larger storm events) which will outlet to an oil/grit separator (OGS) and ultimately to the existing roadside ditch along Borrisokane Road. Permeable pavers for sidewalks are proposed along the western and southern property limits to further reduce runoff from impermeable surfaces.

To prevent any sediment or erosion runoff into the realigned drainage feature, it is recommended that ESC fencing be installed around the entire work area prior to any earth works or soil exposure. This will require daily inspection to ensure fencing is doing an adequate job preventing any loose soil from leaving the Subject Property.

## 8. Policy Conformity

A summary of applicable natural heritage policies and the manner in which the proposed development plan meets their requirements is provided in **Table 1**. With the implementation of the abovementioned mitigation measures, there are no predicted negative impacts to the limited Natural Heritage Features observed adjacent to the Subject Property or their ecological functions.

**Table 1. Policy Conformity**

Policy Document	Policy Intent/Objective	Implications and Policy Conformity
Migratory Bird Act	Protect most species of migratory birds and their nests and eggs anywhere they are found in Canada.	Tree clearing and vegetation removal is not anticipated for the proposal as no suitable vegetation currently exists within the work area.
Endangered Species Act	Species and the habitat of species designated as Endangered or Threatened are afforded legal protection.	The proposed development occurs within anthropogenic and cultural meadow areas. These areas do not have existing structures or appropriate vegetation that may provide habitat (i.e., existing structures, grassland). Risk to SAR species or their habitat are not anticipated. Mitigation (e.g., plantings, ESC fencing) will mitigate indirect impacts into the adjacent watercourse system.
Provincial Policy Statement	Direction to regional and local municipalities regarding planning policies for the protection and management of natural heritage features.	Development or site alteration is not to occur within Significant Woodlands or Wetlands in Ecoregion 6E. None of these features have been identified within or immediately adjacent to the Subject Property. Development is proposed outside of these natural features and their adjacent lands.
City of Ottawa Official Plan (2021)	To manage this growth in ways that reinforce the qualities of the City of Ottawa most valued by its residents (i.e., its distinctly livable communities, its green and open character, and its unique characteristics)	No Natural Heritage Features were found on the Subject Property or on adjacent lands (within 120 m), with the exception of the realigned drainage feature to the east. Watercourses to the west and south of the Subject Property were no longer present. Mitigation (e.g., plantings, ESC fencing) will mitigate indirect impacts into the adjacent watercourse system.  The City of Ottawa confirmed that, as per provision 3 of Section 69 of the zoning by-law, no formal setback from the realigned drainage feature will be required on the Subject Property. However, a planting plan using native species has been provided to merge with this adjacent watercourse area.
Rideau Valley Conservation Authority ( <i>O. Reg. 174/06</i> )	RVCA regulates activities in wetlands, valleylands, watercourses, and hazardous areas.	The Subject Property does contain Regulated Area (as identified by RVCA mapping) or regulated features and therefore consultation and an application of permission will be required for the proposed development.

## 9. Conclusions

The findings of this EIS are the result of a background review, field investigations, and an assessment of ecological data, as well as the current natural heritage policy requirements. We have identified the natural environmental sensitivities, constraints and development opportunities associated with the proposed development plan. Based on the findings and recommendations of this study, the only natural heritage features which requires consideration is that of the realigned watercourse which is within 120 metres of the Subject Property boundaries. Potential negative impacts to this realigned watercourse will be mitigated through:

- ESC measures – Measures outlined in Civil Plan C101 will prevent excess sediments and soils from entering the watercourse.
- Native plantings along the surrounding and eastern boundary of the Subject Property – These plantings will provide shading effects, offsets for vehicular emissions, and erosion prevention.
- Light, air and noise pollution measures – Mitigation measures outlined in section 7.2 will prevent excess light and noise from impacting the watercourse.
- Debris measures - Measures outlined in Site Plan SP01 will provide opportunities for garbage storage. Thus, debris will be prevented from entering the watercourse.
- Adherence to wildlife protection protocols – Through the use of proper garbage receptacles and proper drainage measures, unwanted wildlife will not be attracted to the area. Additionally, the clearing of the vegetation should begin on the west side of the Subject Property and progress east towards the watercourse, giving any potential wildlife the opportunity to move towards the natural feature.

It is our professional opinion that with the implementation of the mitigation measures provided in this report, that the realigned watercourse and its ecological functions will not be negatively impacted by the proposed development plan and adheres to the existing environmental framework.

## 10. Certification

This report was prepared, reviewed, and approved by the undersigned:

**Prepared By:**



---

Hayley Brown, B.Sc.  
Ecologist

**Reviewed By:**



---

Austin Adams, B.Sc.  
Senior Ecologist

# 11. References

- Bird Studies Canada. (2023). *Atlas of the Breeding Birds of Ontario*. Retrieved from Atlas of the Breeding Birds of Ontario: <http://www.birdsontario.org/atlas/index.jsp?lang=en>
- Brunton, D. F. (2005). *Ottawa Natural Areas Environmental Evaluation Study: Appendix A - Vascular Plants of the City of Ottawa with Identification of Significant Species*. Ottawa: City of Ottawa.
- City of Ottawa. (2006). *Jock River Reach 1 Subwatershed Study*. Retrieved from City of Ottawa: <https://app06.ottawa.ca/calendar/ottawa/citycouncil/a-ofac/2006/10-23/10-23/agendaitem9.htm>
- City of Ottawa. (2015). *City of Ottawa Protocol for Wildlife Protection during Construction*. . Retrieved from City of Ottawa: [https://documents.ottawa.ca/sites/documents/files/documents/construction\\_en.pdf](https://documents.ottawa.ca/sites/documents/files/documents/construction_en.pdf)
- City of Ottawa. (2015). *Environmental Impact Statement Guidelines*. Retrieved from City of Ottawa: [https://documents.ottawa.ca/sites/documents/files/documents/eis\\_guidelines2015\\_en.pdf](https://documents.ottawa.ca/sites/documents/files/documents/eis_guidelines2015_en.pdf)
- City of Ottawa. (2021). *City of Ottawa Official Plan*. Retrieved from City of Ottawa: <https://engage.ottawa.ca/the-new-official-plan>
- City of Ottawa. (2021). *Zoning (By-law No. 2008-250)*. Retrieved from City of Ottawa: <https://ottawa.ca/en/living-ottawa/laws-licences-and-permits/laws/laws-z/zoning-law-no-2008-250>
- Government of Canada. (1994). Migratory Birds Convention Act, 1994 (S.C. 1994, c. 22). Retrieved from <http://laws-lois.justice.gc.ca/eng/acts/m-7.01/>
- Government of Canada. (2018). *Beneficial Management Practices - Avoiding Harm to Migratory Birds*. Retrieved from canada.ca: <https://www.canada.ca/en/environment-climate-change/services/avoiding-harm-migratory-birds/beneficial-management-practices.html>
- Government of Ontario. (2007). Endangered Species Act, 2007, S.O. 2007, c. 6. Retrieved from <https://www.ontario.ca/laws/statute/07e06>
- Government of Ontario. (2023). *Land Information Ontario*. Retrieved from ontario.ca: <https://www.ontario.ca/page/land-information-ontario>
- Lee, H. T., Bakowsky, W. D., Riley, J., Bowles, J., Puddister, M., Uhlig, P., & McMurray, S. (1998). *Ecological Land Classification for Southern Ontario: First Approximation and its Application*. Ontario Ministry of Natural Resources, Southcentral Science Section, Science Development and Transfer Branch.



- Ministry of Natural Resources and Forestry. (2018). *Natural Heritage Information Request Guide*. Ministry of Natural Resources and Forestry.
- Ministry of Natural Resources and Forestry. (2023). *Make a Map: Natural Heritage Areas*. Retrieved from Ministry of Natural Resources and Forestry - Make a Map: Natural Heritage Areas: [http://www.gisapplication.lrc.gov.on.ca/mamnh/Index.html?site=MNR\\_NHLUPS\\_NaturalHeritage&viewer=NaturalHeritage&locale=en-US](http://www.gisapplication.lrc.gov.on.ca/mamnh/Index.html?site=MNR_NHLUPS_NaturalHeritage&viewer=NaturalHeritage&locale=en-US)
- Ministry of Natural Resources and Forestry. (2023). *Natural Heritage Information Centre Species Lists*. Retrieved from Ministry of Natural Resources and Forestry: <https://www.ontario.ca/page/get-natural-heritage-information>
- Morton, J. K., & Venn, J. M. (1984). *The Flora of Manitoulin Island. 2nd Ed. Biology Series No. 28*. Waterloo, Ontario: University of Waterloo.
- Oldham, M. J., & Brinker, S. R. (2009). *Rare Vascular Plants of Ontario, Fourth Edition*. Peterborough, Ontario: Natural Heritage Information Centre, Ministry of Natural Resources.
- Oldham, M. J., Bakowsky, W. D., & Sutherland, D. A. (1995). *Floristic Quality Assessment System for Southern Ontario*. Peterborough, ON: Natural Heritage Information Centre, Ontario Ministry of Natural Resources.
- Ontario Ministry of Municipal Affairs and Housing. (2020). *Provincial Policy Statement, 2020*. Toronto, ON. doi:ISBN 978-1-4606-3522-3
- Ontario Ministry of Natural Resources. (2010). *Natural Heritage Reference Manual for Natural Heritage Policies of the Provincial Policy Statement, 2005. Second Edition*. Toronto: Queen's Printer for Ontario. Retrieved from <http://cloca.ca/resources/Outside%20documents/Natural%20Heritage%20Policies%20of%20the%20Provincial%20Policy%20Statement%20MNR%202010.pdf>
- Ontario Ministry of Natural Resources and Forestry. (2015). *Significant Wildlife Habitat Criteria Schedules For Ecoregion 6E*. Peterborough: Regional Operations Division, Southern Region Resources Section.
- Ontario Nature. (2023). *Ontario Reptile and Amphibian Atlas*. Retrieved from Ontario Nature: <https://ontarionature.org/programs/citizen-science/reptile-amphibian-atlas/>

Rhuland and Associates Ltd. (2023). *Landscape Plan L-01 - 652 Flagstaff Drive*. Ottawa.

Rideau Valley Conservation Authority. (2013). *Rideau Valley Conservation Authority: Regulation of Development, Interference with Wetlands and Alterations to Shorelines and Watercourses, O Reg 174/06*. Retrieved from Rideau Valley Conservation Authority:  
[https://www.rvca.ca/media/k2/attachments/13elaws\\_regs\\_060Z174\\_eZRVCA\\_1.pdf](https://www.rvca.ca/media/k2/attachments/13elaws_regs_060Z174_eZRVCA_1.pdf)



# Appendix A

## Species at Risk Screening

## Appendix A: Species at Risk Screening

NAME	SARA STATUS	SARO	COSEWIC	SCHEDULE	S-RANK	HABITAT REQUIREMENTS	SOURCE OF RECORD	POTENTIAL HABITAT PRESENT (Y/N)	RATIONALE	POTENTIAL IMPACTS AND MITIGATION
<b>AVIFAUNA</b>										
Bank Swallow ( <i>Riparia riparia</i> )	THR	THR	THR	1	S4B	The Bank Swallow is threatened by loss of breeding and foraging habitat, destruction of nesting habitat and widespread pesticide use. Bank swallows are small songbirds with brown upperparts, white underparts and a distinctive dark breast band. It averages 12 cm long and weighs between 10 and 18 grams. The swallow can be distinguished in flight from other swallows by its quick, erratic wing beats and its almost constant buzzy, chattering vocalizations. They nest in burrows in natural and human-made settings where there are vertical faces in silt and sand deposit, including banks of rivers and lakes, active sand and gravel pits or former ones where the banks remain suitable. The birds breed in colonies ranging from several to a few thousand pairs (Ministry of Natural Resources and Forestry, 2014).	OBBA	N	No habitat available.	N/A
Barn Swallow ( <i>Hirundo rustica</i> )	THR	THR	THR	1	S4B	The Barn Swallow is a threatened species, is found throughout southern Ontario, and can range into the north as long as suitable nesting locations can be found. These birds prefer to nest within human made structures such as barns, bridges, and culverts. Barn Swallow nests are cup-shaped and made of mud; they are typically attached to horizontal beams or vertical walls underneath an overhang. A significant decline in populations of this species has been documented since the mid-1980s, which is thought to be related to a decline in prey. Since the Barn Swallow is an aerial insectivore, this species relies on the presence of flying insects at specific times during the year. Changes in building practices and materials may also be having an impact on this species (Ministry of Natural Resources and Forestry, 2015).	OBBA	N	No habitat available.	N/A
Bobolink ( <i>Dolichonyx oryzivorus</i> )	THR	THR	THR	1	S4B	The Bobolink is found in grasslands and hayfields, and feeds and nests on the ground. This species is widely distributed across most of Ontario; however, are designated at risk because of rapid population decline over the last 50 years (Ministry of Natural Resources and Forestry, 2014). The historical habitat of the bobolink was tallgrass prairie and other natural open meadow communities; however, as a result of the clearing of native prairies and the post-colonial increase in agriculture, bobolinks are now widely found in hayfields. Due to their reproductive cycle, nesting habits, and use of agricultural areas, bobolink nests and young are particularly vulnerable to loss as a result of common agricultural practices (i.e. first cut hay).	OBBA and NHIC	N	No grassland habitat available.	Bobolink prefer open grasslands which can include Cultural Meadows (CUM) as described on the property. This species requires grassland meadow habitat which would have been evidenced by persistent grass stalks. The meadow within the property is a sparsely regenerating forb meadow.
Eastern Meadowlark ( <i>Sturnella magna</i> )	THR	THR	THR	1	S4B	The Eastern Meadowlark is a bird that prefers pastures and hayfields, but is also found to breed in orchards, shrubby fields and human use areas such as airports and roadsides. Eastern meadowlarks can nest from early May to mid-August, in nests that are built on the ground and well-camouflaged with a roof woven from grasses. The decline in population of these species is thought to be at least partially related to habitat destruction and agricultural practices (Ministry of Natural Resources and Forestry, 2014).	OBBA	N	No grassland habitat available.	Eastern Meadowlark also prefer open grasslands which can include Cultural Meadows (CUM) as described on the property. This species requires grassland meadow habitat which would have been evidenced by persistent grass stalks. The meadow within the property is a sparsely regenerating forb dominated meadow.
Eastern Wood-Pewee ( <i>Contopus virens</i> )	SC	SC	SC	1	S4B	The Eastern Wood-pewee is classified as a species of special concern by COSSARO. Their population has been gradually declining since the mid-1960's (The Cornell Lab of Ornithology, 2015). The Eastern Wood-pewee is a "flycatcher", a bird that eats flying insects, that lives in the mid-canopy layer of forest clearings and edges of deciduous and mixed forests. It prefers intermediate-age forest stands with little understory vegetation. Threats to the population are largely unknown; however, causes may include loss of habitat due to urban development and decreases in the availability of flying insect prey (Ministry of Natural Resources and Forestry, 2014).	OBBA	N	Woodlands not found within property	N/A
Grasshopper Sparrow ( <i>Ammodramus savannarum</i> )	No Status	No Status	SC	X	S4B	Grasshopper Sparrow are specialized to open relatively short grassland habitat, preferably grasslands with relatively sparse cover such as those in areas of poor soils, including alvars, moraines, and sand plains and generally does not favour tall grass moist meadows. It will also breed in manmade hayfields and occasionally in cereals such as Rye ( <i>Secale cereale</i> ).	OBBA	N	No grassland habitat available.	N/A
Wood Thrush ( <i>Hylocichla mustelina</i> )	THR	SC	THR	1	S4B	The Wood Thrush is a species of Special Concern because of habitat degradation or destruction by anthropogenic development. The Wood Thrush is a medium-sized songbird, generally rusty-brown on the upper parts with white under parts and large blackish spots on the breast and sides, and about 20 cm long. The Wood Thrush forages for food in leaf litter or on semi-bare ground, including larval and adult insects as well as plant material. They seek moist stands of trees with well-developed undergrowth in large mature deciduous and mixed (conifer-deciduous) forests. The Wood Thrush flies south to Mexico and Central America for the winter (Ministry of Natural Resources and Forestry, 2014).	OBBA	N	Woodlands not found within property	N/A
<b>HERPTILES</b>										

NAME	SARA STATUS	SARO	COSEWIC	SCHEDULE	S-RANK	HABITAT REQUIREMENTS	SOURCE OF RECORD	POTENTIAL HABITAT PRESENT (Y/N)	RATIONALE	POTENTIAL IMPACTS AND MITIGATION
Blanding's Turtle ( <i>Emydoidea blandingii</i> )	THR	THR	END	1	S3	Blanding's turtles are threatened in Ontario primarily as a result of habitat loss and fragmentation. Blanding's turtles spend the majority of their life cycle in the aquatic environment, using terrestrial sites for travel between habitat patches and to lay clutches of eggs. These turtles prefer shallow nutrient rich water with organic sediment and dense vegetation. Blanding's turtles nest in dry coniferous and mixed forest habitats, as well as fields and roadsides (Government of Canada, 2015).	ORAA	N	No habitat available within property	N/A
Northern Map Turtle ( <i>Graptemys geographica</i> )	SC	SC	SC	1	S3	The northern map turtle is a medium sized turtle with a carapace marked by concentric rings that resemble contour lines on a map. The range of this turtle includes larger lakes and rivers that contain an abundance of their primary prey species; molluscs. Shoreline development, water pollution and the spread of the zebra mussel are notable reasons for the decline in populations of this species (Ministry of Natural Resources and Forestry, 2014).	ORAA	N	No habitat available within property	N/A
Snapping Turtle ( <i>Chelydra serpentina</i> )	SC	SC	SC	1	S3	The snapping turtle is a species of special concern in Ontario due to the potential for the species to become threatened or endangered as a result of biological factors or other identified threats. While not presently protected by law, the snapping turtle has been recognized as a species of special concern by COSSARO. Snapping turtles spend the majority of their lives in water and travel slightly upland to gravel or sandy embankments or beaches to lay their eggs (Ontario Ministry of Natural Resources and Forestry, 2014).	ORAA	N	No habitat available within property	N/A
<b>VASCULAR PLANTS</b>										
Butternut ( <i>Juglans cinerea</i> )	END	END	END	1	S2?	The butternut is designated as endangered by COSSARO and is tracked by the NHIC as a species at risk. The tree is federally regulated by the Species at Risk Act (2002). Butternut belongs to the walnut family and produces edible nuts which are a preferred food source for wildlife. The range of butternut trees is south of the Canadian Shield on soils derived from calcium rich limestone bedrock. Butternut trees, which at one time were much more common to the south extending to the northern aspect of zone 6E, have been declining due to factors including forest loss and disease. Butternut trees suffer from a highly transmissible fungal disease called butternut canker. Butternut canker is causing very rapid decline in this tree species across its native range. The fungal disease is easily transmitted by wind and is very difficult to prevent. Trees often die within a few years of infection by butternut canker (Ministry of Natural Resource and Forestry, 2014).	NHIC	N	No woodland or treed areas located on or immediately adjacent to the property.	No Butternut trees were detected on the property.
<b>MAMMALS</b>										
Tri-colored Bat (Eastern Pipistrelle) ( <i>Perimyotis subflavus</i> )	END	END	END	1	S3?	The eastern pipistrelle is a small bat that is widely distributed in eastern North America and whose range extends north to southern Ontario. The eastern pipistrelle is rare in this region of Ontario which is at the northernmost limit of the natural range for the species. These bats prefer to nest in foliage, tree cavities and woodpecker holes, and are occasionally found in buildings; though this is not their preferred habitat. Winter hibernation takes place in caves, mines and deep crevices. Eastern pipistrelles feed primarily on small insects and prefer an open forest habitat type in proximity to water (University of Michigan Museum of Zoology, 2004).	Professional Experience	N	No treed habitats on property	N/A
Eastern Small-footed Myotis ( <i>Myotis leibii</i> )	No Status	END	No Status	No Scheduling	S2S3	The eastern small-footed myotis, a bat, are an endangered species threatened by a disease known as white nose syndrome, caused by a fungus from Europe. Eastern small-footed bat's fur has black roots and shiny light brown tips, giving it a yellowish-brown appearance. Its face mask, ears and wings are black, and its underside is grayish-brown, about 8 cm long in size and weighs 4-5 grams. In the spring and summer, eastern small-footed bats will roost in a variety of habitats, including in or under rocks, in rock outcrops, in buildings, under bridges, or in caves, mines, or hollow trees. They change their roosting locations daily and hunt at night for insects to eat, including beetles, mosquitos, moths, and flies. They hibernate in winter, often in caves and abandoned mines. They can be found from south of Georgian Bay to Lake Erie and east to the Pembroke area, and choose colder and drier sites (Ministry of Natural Resources and Forestry, 2014).	Professional Experience	N	No treed habitats on property	N/A
Little Brown Myotis ( <i>Myotis lucifugus</i> )	END	END	END	1	S4	Little brown myotis, a bat, are an endangered species threatened by a disease known as white nose syndrome, caused by a fungus from Europe. Little brown bats have glossy brown fur and usually weigh between four and 11 grams. Bats are nocturnal. During the day they roost in trees and buildings. They often select attics, abandoned buildings and barns for summer colonies where they can raise their young. Little brown bats hibernate from October or November to March or April, most often in caves or abandoned mines that are humid and remain above freezing – an ideal environment for the fungus to grow and flourish. The syndrome affects bats by disrupting their hibernation cycle, so that they use up body fat supplies before the spring when they can once again find food sources (Ministry of Natural Resources and Forestry, 2014).	Professional Experience	N	No treed habitats on property	N/A

NAME	SARA STATUS	SARO	COSEWIC	SCHEDULE	S-RANK	HABITAT REQUIREMENTS	SOURCE OF RECORD	POTENTIAL HABITAT PRESENT (Y/N)	RATIONALE	POTENTIAL IMPACTS AND MITIGATION
Northern Myotis ( <i>Myotis septentrionalis</i> )	END	END	END	1	S3	The northern long-eared myotis, a bat, are an endangered species threatened by a disease known as white nose syndrome, caused by a fungus from Europe. Northern long-eared bats have dull yellow-brown fur with pale grey bellies. They are approximately eight cm long, with a wingspan of about 25 cm, and usually weigh six to nine grams. Northern long-eared bats can be found in boreal forests, roosting under loose bark and in the cavities of trees. These bats hibernate from October or November to March or April, most often in caves or abandoned mines (Ministry of Natural Resources and Forestry, 2014).	Professional Experience	N	No treed habitats on property	N/A
<b>OTHER</b>										
Monarch Butterfly ( <i>Danaus plexippus</i> )	SC	SC	END	1	S2N,S4B	The monarch is an orange and black butterfly with small white spots and is classified as a species of special concern by COSSARO. The monarch relies on milkweed plants as a food source for growing caterpillars, but the adult butterflies forage in diverse habitats for nectar from wildflowers. The greatest threat to the monarch is loss of overwintering habitat in Mexico. Other threats include use of pesticides and herbicides throughout its range (Ministry of Natural Resources and Forestry, 2014).	OBA	N	Suitable meadow habitat not present	N/A

**Notes:**

- SC - Special Concern
- THR - Threatened
- END - Endangered
- S1 - Extremely rare in Ontario
- S2 - Very rare in Ontario
- S3 - Rare to uncommon in Ontario
- S4 - Considered to be common in Ontario
- S5 - Species is widespread in Ontario
- SH - Possibly extirpated
- S#S# - Indicates insufficient information exists to assign a single rank.
- S#? - Indicates some uncertainty with the classification due to insufficient data.
- S#N - Nonbreeding
- S#B - Breeding



# Appendix B

## Plant List

## Appendix B – Flora List

Family	Scientific Name	Common Name	COSEWIC	SARA	SARO	NHIC Status			Coefficient of Conservatism	Coefficient of Wetness	Regional Rarity Ottawa
						Global	Provincial	Exotic			
Poaceae	<i>Agrostis gigantea</i>	Redtop				G4G5	NNA	SE5		-3	Common
Brassicaceae	<i>Alliaria petiolata</i>	Garlic Mustard				GNR	NNA	SE5		0	Common
Asteraceae	<i>Arctium minus</i>	Common Burdock				GNR	NNA	SE5		3	Common
Asteraceae	<i>Artemisia vulgaris</i>	Common Wormwood				GU	NNA	SE5		5	Uncommon (local along Ottawa River)
Brassicaceae	<i>Barbarea vulgaris</i>	Bitter Wintercress				GNR	NNA	SE5		0	Common
Poaceae	<i>Bromus inermis</i>	Smooth Brome				G5	NNA	SE5		5	Common
Cyperaceae	<i>Carex stipata</i>	Awl-fruited Sedge				G5	N5		3	-5	Common
Asteraceae	<i>Cirsium vulgare</i>	Bull Thistle				GNR	NNA	SE5		3	Common
Apiaceae	<i>Daucus carota</i>	Wild Carrot				GNR	NNA	SE5		5	Common
Equisetaceae	<i>Equisetum arvense</i>	Field Horsetail				G5	N5		0	0	Common
Asteraceae	<i>Erigeron annuus</i>	Annual Fleabane				G5	N5		0	3	Common
Asteraceae	<i>Erigeron canadensis</i>	Canada Horseweed				G5	N5		0	3	Common
Asteraceae	<i>Lactuca sp.</i>	Lettuce Species									Common
Asteraceae	<i>Leucanthemum vulgare</i>	Oxeye Daisy				GNR	NNA	SE5		5	Common
Poaceae	<i>Lolium sp.</i>	Ryegrass Species									Common
Fabaceae	<i>Lotus corniculatus</i>	Garden Bird's-foot Trefoil				GNR	NNA	SE5		3	Common
Apiaceae	<i>Pastinaca sativa</i>	Wild Parsnip				GNR	NNA	SE5		5	Common

**Legend:**

COSEWIC - Committee on the Status of Endangered Wildlife in Canada

NHIC - Natural Heritage Information Centre

SARA - Ontario Species at Risk Act List

Coefficients (Conservatism and Wetness) – Oldham et al., 1995

SARO - Species at Risk in Ontario

Ottawa Rank – Brunton, 2005



Family	Scientific Name	Common Name	COSEWIC	SARA	SARO	NHIC Status			Coefficient of Conservatism	Coefficient of Wetness	Regional Rarity Ottawa
						Global	Provincial	Exotic			
Poaceae	<i>Phalaris arundinacea</i>	Reed Canarygrass				G5	N5		0	-3	Common
Plantaginaceae	<i>Plantago major</i>	Common Plantain				G5	NNA	SE5		3	Common
Poaceae	<i>Poa pratensis</i>	Kentucky Bluegrass				G5	N5		0	3	Common
Brassicaceae	<i>Sinapis arvensis</i>	Corn Mustard				GNR	NNA	SE5		5	Common
Asteraceae	<i>Solidago canadensis</i>	Canada Goldenrod				G5	N5		1	3	Common
Asteraceae	<i>Symphotrichum novae-angliae</i>	New England Aster				G5	N5		2	-3	Common
Asteraceae	<i>Taraxacum officinale</i>	Common Dandelion				G5	N5	SE5		3	Common
Brassicaceae	<i>Thlaspi arvense</i>	Field Pennycress				GNR	NNA	SE5		5	Common
Fabaceae	<i>Trifolium hybridum</i>	Alsike Clover				GNR	NNA	SE5		3	Common
Fabaceae	<i>Trifolium pratense</i>	Red Clover				GNR	NNA	SE5		3	Common
Fabaceae	<i>Trifolium repens</i>	White Clover				GNR	NNA	SE5		3	Common
Asteraceae	<i>Tussilago farfara</i>	Coltsfoot				GNR	NNA	SE5		3	Common
Scrophulariaceae	<i>Verbascum thapsus</i>	Common Mullein				GNR	NNA	SE5		5	Common
Scrophulariaceae	<i>Veronica sp.</i>	Speedwell Species									Common
Fabaceae	<i>Vicia cracca</i>	Tufted Vetch				GNR	NNA	SE5		5	Common

**Legend:**

COSEWIC - Committee on the Status of Endangered Wildlife in Canada

SARA - Ontario Species at Risk Act List

SARO - Species at Risk in Ontario

NHIC - Natural Heritage Information Centre

Coefficients (Conservatism and Wetness) – Oldham et al., 1995

Ottawa Rank – Brunton, 2005



# Appendix C

## **Significant Wildlife Habitat Screening**

SWH Type	Associated Species	Associated ELC Ecosites	Habitat Criteria	Presence (Y/N)	Additional Notes and Species Observations
<b>Seasonal Concentration Areas of Animals</b>					
Waterfowl Stopover and Staging Areas (Terrestrial)	Ducks	CUM + CUT ecosites	Fields with sheet-water flooding mid-March to May	N	No sheet flooding of field identified within property
Waterfowl Stopover and Staging Area (Aquatic)	Ducks, Geese	Ponds, Lakes, Inlets, Marshes, Swamps, Shallow Water Ecosites	Sewage & SWM ponds <b>not</b> SWH. Reservoir managed as a large wetland or pond/lake qualifies.	N	Habitat not found within property.
Shorebird Migratory Stopover Area	Shorebirds	Beaches, Dunes, Meadow Marshes	Shorelines. Sewage treatment ponds and storm water ponds <b>not</b> SWH.	N	Habitat not found within property.
Raptor Wintering Area	Eagles, Hawks, Owls	<b>Hawks/Owls:</b> Combination of both Forest and Cultural Ecosites <b>Bald Eagle:</b> Forest or swamp near open water (hunting ground)	<b>Raptors:</b> >20ha, with a combo of forest and upland. Meadow (>15ha) with adjacent woodlands. <b>Eagles:</b> open water, large trees & snags for roosting.	N	Habitat not found within property.
Bat Hibernacula	Big Brown Bat, Tri-coloured Bat	Caves, Crevices, mines, karsts	Buildings and active mine sites <b>not</b> SWH.	N	Habitat not found within property.
Bat Maternity Colonies	Big Brown Bat, Silver-haired Bat	Deciduous or mixed forests and swamps.	Mature deciduous and mixed forests with >10/ha cavity trees >25 cm DBH.	N	Habitat not found within property.
Turtle Wintering Area	<b>Turtles</b> (Midland, N. Map, Snapping)	SW, MA, OA, SA, FEO, BOO (requires open waters)	<b>Free water beneath ice.</b> Soft mud substrate. Permanent water bodies, large wetlands, bogs, fens with adequate DO.	N	Habitat not found within property.
Reptile Hibernaculum	Snakes	<b>Snakes:</b> Any ecosite (esp. w/ rocky areas), other than very wet ones. <b>Five-lined Skink:</b> FOD and FOM, FOC1, FOC3 - with rock outcrops	<b>Access below frost line:</b> burrows; <b>rock</b> crevices, piles or slopes, <b>stone</b> fences or foundations. Conifer/shrubby swamps/swales, poor fens, depressions in bedrock w/ accumulations of sphagnum moss or sedge hummock ground cover.	N	Habitat not found within property.
Colonially-nesting Bird Breeding Habitat (Bank and Cliff)	Cliff Swallow, N. Rough-winged Swallow	Banks, sandy hills/piles, pits, slopes, cliff faces, bridge abutments, silos, barns.	Exposed soil banks, <b>not</b> a licensed/permitted aggregate area or new man-made features (2 yrs).	N	Habitat not found within property.
Colonially-nesting Bird Breeding Habitat (Tree/Shrubs)	Great Blue Heron, Black-crowned NightHeron, Great Egret, Green Heron	SWM2, SWM3, SWM5, SWM6, SWD1 to SWD7, FET1	Nests in live or dead standing trees in wetlands, lakes, islands and peninsulas. Shrubs and emergents may be used. Nests in trees are 11 - 15 m from ground, near tree tops.	N	Habitat not found within property.
Colonially-nesting Bird Breeding Habitat (Ground)	Herring Gull, Great Black-backed Gull, Little Gull, Ring-billed Gull, Common Tern, Caspian Tern, Brewer's Blackbird	<b>Gulls/Terns:</b> Rocky island or peninsula in lake or river. <b>Brewer's Blackbird:</b> close to watercourses in open fields or pastures with scattered trees or shrubs.	<b>Gulls/Terns:</b> islands or peninsulas with open water or marshy areas. <b>Brewers Blackbird colonies:</b> on the ground in low bushes close to streams and irrigation ditches.	N	Habitat not found within property.
Migratory Butterfly Stopover Area	Painted Lady, Red Admiral, <b>Special Concern:</b> Monarch	Combination of open (CU) and forested (FO) ecosites (need one from each).	≥10 ha, located within 5 km of Lake Ontario. Undisturbed sites, with preferred nectar species.	N	Habitat not found within property.
Landbird Migratory Stopover Areas	All migratory songbirds. All migrant raptor species.	Forest (FO) and Swamp (SW) ecosites	Woodlots >10 ha within 5 km of Lake Ontario. If multiple woodlands are along the shoreline, those <2 km from L. Ontario are more significant.	N	Habitat not found within property.
Deer Yarding Areas	White-tailed Deer	Mixed or Conifer ecosites	Determined by MNRF - no studies	N	Habitat not found within property.
Deer Winter Congregation Areas	White-tailed Deer	Mixed or Conifer ecosites	Determined by MNRF - no studies	N	Habitat not found within property.
<b>Rare Vegetation Communities</b>					
Cliffs and Talus Slopes		TAO, TAS, CLO, CLS, TAT, CLT e.g., Niagara Escarpment (contact NEC)	<b>Cliff:</b> near vertical bedrock >3m <b>Talus Slope:</b> coarse rock rubble at the base of a cliff	N	Habitat not found within property.
Sand Barren		SBO1, SBS1, SBT1	Sand Barrens >0.5 ha. Vegetation can vary from patchy and barren to tree covered, but <60%. <50% vegetation cover are exotic species.	N	Habitat not found within property.
Alvar	<i>Carex crawei</i> , <i>Panicum philadelphicum</i> , <i>Eleocharis compressa</i> , <i>Scutellaria parvula</i> , <i>Trichostema brachiatum</i> , Loggerhead Shrike	ALO1, ALS1, ALT1, FOC1, FOC2, CUM2, CUS2, CUT2-1, CUW2	Alvar >0.5 ha. <b>Need 4 of the 5 Alvar Indicator Spp.</b> <50% vegetation cover are exotic species.	N	Habitat not found within property.
Old Growth Forest	Trees >140 yrs; heavy mortality = gaps. Multi-layer canopy, lots of snags and downed logs	FOD, FOC, FOM, SWD, SWC, SWM	Woodland areas ≥30 ha with ≥10 ha interior habitat, assuming a 100 m buffer at edge of forest.	N	Habitat not found within property.
Savannah	Prairie Grasses w/ trees	TPS1, TPS2, TPW1, TPW2, CUS2	A Savannah is a <b>tallgrass prairie</b> habitat that has tree cover of 25 – 60%. <50% cover of exotic species.	N	Habitat not found within property.
Tallgrass Prairie	Prairies Grasses dominate	TPO1, TPO2	An <b>open Tallgrass Prairie</b> habitat has < 25% tree cover. Less than 50% cover of exotic species.	N	Habitat not found within property.
Other Rare Vegetation Communities		Provincially Rare S1 - S3 veg. comm. are listed in Appendix M of SWHTG.	Rare Vegetation Communities may include beaches, fens, forest, marsh, barrens, dunes and swamps.	N	Habitat not found within property.
<b>Specialized Habitat for Wildlife</b>					

SWH Type	Associated Species	Associated ELC Ecosites	Habitat Criteria	Presence (Y/N)	Additional Notes and Species Observations
Waterfowl Nesting Area	Ducks	Upland habitats adjacent to: MAS1 to MAS3, SAS1, SAM1, SAF1, MAM1 to MAM6, SWT1, SWT2, SWD1 to SWD4 (>0.5 ha open water wetlands, alone or collectively).	Extends 120 m from a wetland or wetland complex. Upland areas should be at least 120 m wide. Wood Ducks and Hooded Mergansers use cavity trees (>40 cm dbh).	N	Habitat not found within property.
Bald Eagle & Osprey Nesting, Foraging and Perching Habitat	Osprey, Bald Eagle	FOD, FOM, FOC, SWD, SWM, SWC directly adjacent to riparian areas	Nesting areas are associated with waterbodies along forested shorelines, islands, or on structures over water.	N	Habitat not found within property.
Woodland Raptor Nesting Habitat	Barred Owl. <b>Hawks:</b> N. Goshawk, Cooper's, Sharp-shinned, Red-shouldered, Broad-winged.	Forests (FO), swamps (SW), and conifer plantations	>30 ha with > 10 ha interior habitat.	N	Habitat not found within property.
Turtle Nesting Areas	Midland Painted Turtle <b>Special Concern:</b> Snapping Turtle, Northern Map Turtle	Exposed mineral soil (sand or gravel) areas adjacent (<100m) or within: MAS1 to MAS3, SAS1, SAM1, SAF1, BOO1	Nest sites within open sunny areas with soil suitable for digging. Sand and gravel beaches.	N	Habitat not found within property.
Seeps and Springs	Wild Turkey, Ruffed Grouse, Spruce Grouse, White-tailed Deer, Salamander spp.	Seeps/Springs are areas where ground water comes to the surface.	Any forested area within the headwaters of a stream/river system. <b>(2 or more confirms SWH type).</b>	N	Habitat not found within property.
Amphibian Breeding Habitat (Woodland)	Woodland Frogs and Salamanders	FOC, FOM, FOD, SWC, SWM, SWD	Open water wetlands, pond or woodland pool of >500 m <sup>2</sup> within or adjacent to wooded areas. Permanent ponds or holding water until mid-July preferred.	N	Habitat not found within property.
Amphibian Breeding Habitat (Wetlands)	Toads, Frogs, and Salamanders	SW, MA, FE, BO, OA and SA. Typically isolated (>120m) from woodland ecosites, however larger wetlands may be adjacent to woodlands.	Open water wetland ecosites >500m <sup>2</sup> isolated from woodland ecosites with high species diversity. Permanent water with abundant vegetation for bullfrogs.	N	Habitat not found within property.
Woodland Area-Sensitive Bird Breeding Habitat	Birds (area-sensitive species)	FOC, FOM, FOD, SWC, SWM, SWD	Large mature (>60 years) forest stands/woodlots >30 ha. Interior forest habitat >200m from forest edge.	N	Habitat not found within property.
<b>Habitats of Species of Conservation Concern</b>					
Marsh Bird Breeding Habitat	Wetland Birds	MAM1 to MAM6, SAS1, SAM1, SAF1, FEO1, BOO1 <b>Green Heron:</b> SW, MA and CUM1	Wetlands with shallow water and emergent vegetation. Gr. Heron @ edges of these types w/ woody cover.	N	Habitat not found within property.
Open Country Bird Breeding Habitat	Upland Sandpiper, Grasshopper Sparrow, Vesper Sparrow, N. Harrier, Savannah Sparrow, <b>Short-eared Owl (SC)</b>	CUM1, CUM2	Grassland/meadow >30 ha. Not being actively used for farming. Habitat established for 5 years or more.	N	Habitat not found within property.
Shrub/Early Successional Bird Breeding Habitat	<b>Brown Thrasher + Clay-coloured Sparrow (indicators)</b> , Field Sparrow, Black-billed Cuckoo, E. Towhee, Willow Flycatcher, Yellow-breasted Chat, Golden-winged Warbler	CUT1, CUT2, CUS1, CUS2, CUW1, CUW2	Large field areas succeeding to shrub and thicket habitats > 10 ha. Areas not actively used for farming in the last 5 years.	N	Habitat not found within property.
Terrestrial Crayfish	Chimney or Digger Crayfish; Devil Crayfish or Meadow Crayfish	MAM1 to MAM6, MAS1 to MAS3, SWD, SWT, SWM. CUM1 sites with inclusions of the aforementioned.	Wet meadow and edges of shallow marshes (no minimum size) should be surveyed for terrestrial crayfish (typc. protected by wetland setbacks).	N	Habitat not found within property.
Special Concern and Rare Wildlife Species	Any species of concern or rare wildlife species	Any ELC code.	Presence of species of concern or rare wildlife species.	N	Species at Risk discussed in Appendix A and Section 5.1 of report.
<b>Animal Movement Corridors</b>					
Amphibians	Amphibians	all ecosites assoc. w/ water	When Breeding Habitat - wetland confirmed	N	Habitat not found within property.
Deer Movement	White-tailed Deer	all forested ecosites	When Deer Wintering Habitat confirmed	N	Habitat not found within property.
<b>Exceptions for Ecoregion 6E</b>					
Mast Producing: 6E-14	Black Bear	Forested Ecosites	>30 ha w/ mast producing species: Cherry (berries), Oak, Beech (nuts).	N	Habitat not found within property.
Leks: 6E-17	Sharp-tailed Grouse	CUM, CUS, CUT	Grassland/meadow >15 ha adjacent to shrublands, >30 ha adjacent to woodlands. Low agricultural intensity.	N	Cultural Upland Meadow present but Sharp-tailed grouse Lek is an obvious feature and has not been reported in the area.